

Geometric quadratic Chabauty

Bas Edixhoven & Guido Lido

edix@math.leidenuniv.nl guidomaria.lido@gmail.com Math. Inst., Universiteit Leiden

August 25, 2020

Abstract

Since Faltings proved Mordell’s conjecture (1983) we know that the sets of rational points on curves of genus at least 2 are finite. Determining these sets, in individual cases, is still an unsolved problem. Chabauty’s method (1941) is to intersect, for a prime number p , in the p -adic Lie group of p -adic points of the jacobian, the closure of the Mordell-Weil group with the p -adic points of the curve. If the Mordell-Weil rank is less than the genus then this method has never failed.

Minhyong Kim’s non-abelian Chabauty programme aims to remove the condition on the rank. The simplest case, called quadratic Chabauty, was developed by Balakrishnan, Besser, Dogra, Müller, Tuitman and Vonk, and applied in a tour de force to the so-called cursed curve (rank and genus both 3).

This article aims to make the quadratic Chabauty method *small* and *geometric* again, by describing it in terms of only ‘simple algebraic geometry’ (line bundles over the jacobian and models over the integers).

Keywords: rational points, curves, Chabauty, non-abelian, quadratic, geometric.

2010 Mathematical Subject Classification: 14G05, 11G30.

Contents

1	Introduction	2
2	Algebraic geometry	3
3	From algebraic geometry to formal geometry	6
4	Integral points, closure and finiteness	7
5	Parametrisation of integral points, and power series	10
5.1	Logarithm and exponential	11
5.2	Parametrisation by power series	12
5.3	The p -adic closure	14

6	Explicit description of the Poincaré torsor	14
6.1	Norms	14
6.2	Norms along finite relative Cartier divisors	15
6.3	Explicit description of the Poincaré torsor of a smooth curve	16
6.4	Explicit isomorphism for norms along equivalent divisors	19
6.5	Symmetry of the Norm for divisors on smooth curves	20
6.6	Explicit residue disks and partial group laws	22
6.7	Extension of the Poincaré biextension over Néron models	25
6.8	Explicit description of the extended Poincaré bundle	25
6.9	Integral points of the extended Poincaré torsor	28
7	Description of the map from the curve to the torsor	30
8	An example with genus 2, rank 2, and 14 points	31
8.1	The torsor on the jacobian	32
8.2	Some integral points on the biextension	34
8.3	Some residue disks of the biextension	35
8.4	Geometry mod p of integral points	36
8.5	The rational points with a specific image mod 5.	38
8.6	Determination of all rational points	39
9	Some further remarks	40
9.1	Complex uniformisations of some of the objects involved	40
9.2	Finiteness of rational points	42
9.3	The relation with p -adic heights	46
9.4	Work in progress, and future plans	47
	Author contributions	48
	Acknowledgements	48
	References	49

1 Introduction

Faltings proved in 1983, see [13], that for every number field K and every curve C over K of genus at least 2, the set of K -rational points $C(K)$ is finite. However, determining $C(K)$, in individual cases, is still an unsolved problem. For simplicity, we restrict ourselves in this article to the case $K = \mathbb{Q}$.

Chabauty's method (1941) for determining $C(\mathbb{Q})$ is to intersect, for a prime number p , in the p -adic Lie group of p -adic points of the jacobian, the closure of the Mordell-Weil group with the p -adic points of the curve. If the Mordell-Weil rank r satisfies $r < g$ then this method has never failed.

Minhyong Kim’s non-abelian Chabauty programme aims to remove the condition that $r < g$. The ‘non-abelian’ refers to fundamental groups; the fundamental group of the jacobian of a curve is the abelianised fundamental group of the curve. The most striking result in this direction is the so-called quadratic Chabauty method, applied in [5], a technical tour de force, to the so-called cursed curve ($r = g = 3$). For more details we recommend the introduction of [5].

This article aims to make the quadratic Chabauty method *small* and *geometric* again, by describing it in terms of only ‘simple algebraic geometry’ (line bundles over the jacobian, models over the integers, and biextension structures). The main result is Theorem 4.12. It gives a criterion for a given list of rational points to be complete, in terms of points with values in $\mathbb{Z}/p^2\mathbb{Z}$ only. Section 2 describes the geometric method in less than 3 pages, Sections 3–5 give the necessary theory, Sections 6–7 give descriptions that are suitable for computer calculations, and Section 8 treats an example with $r = g = 2$ and 14 rational points. As explained in the remarks following Theorem 4.12, we expect that this approach will make it possible to treat many more curves. Section 9.1 gives some remarks on the fundamental groups of the objects we use. They are subgroups of higher dimensional Heisenberg groups, where the commutator pairing is the intersection pairing of the fundamental group of the curve. Section 9.2 reproves the finiteness of $C(\mathbb{Q})$, for C with $r < g + \rho - 1$, with ρ the rank of the \mathbb{Z} -module of symmetric endomorphisms of the jacobian of C . It also shows that a version of Theorem 4.12 that uses higher p -adic precision will always give a finite upper bound for $C(\mathbb{Q})$. Section 9.3 gives, through an appropriate choice of coordinates that split the Poincaré biextension, the relation between our geometric approach and the p -adic heights used in the cohomological approach. Section 9.4 describes some work in progress, on the extension of our results to arbitrary number fields, applications to modular curves, relation with Arakelov theory, and some plans for the future.

Already for the case of classical Chabauty (working with J instead of T , and under the assumption that $r < g$), where everything is linear, the criterion of Theorem 4.12 can be useful; this has been worked out and implemented by Pim Spelier in his MSc thesis, [25], defended on 2020/06/23. We recommend this work as a gentle introduction into the geometric approach taken in this article. Although this article is about geometry, it contains no pictures. Fortunately, many pictures can be found in [15], and Edixhoven’s lectures at the Arizona Winter School 2020 are online ([12]).

2 Algebraic geometry

Let C be a scheme over \mathbb{Z} , proper, flat, regular, with $C_{\mathbb{Q}}$ of dimension one and geometrically connected. Let n be in $\mathbb{Z}_{\geq 1}$ such that the restriction of C to $\mathbb{Z}[1/n]$ is smooth. Let g be the genus of $C_{\mathbb{Q}}$. We assume that $g \geq 2$ and that we have a rational point $b \in C(\mathbb{Q})$; it extends uniquely to a $b \in C(\mathbb{Z})$. We let J be the Néron model over \mathbb{Z} of the jacobian $\text{Pic}_{C_{\mathbb{Q}}/\mathbb{Q}}^0$. We denote by J^{\vee} the Néron model over \mathbb{Z} of the dual $J_{\mathbb{Q}}^{\vee}$ of $J_{\mathbb{Q}}$, and $\lambda: J \rightarrow J^{\vee}$ the isomorphism extending the canonical principal polarisation of $J_{\mathbb{Q}}$. We let $P_{\mathbb{Q}}$ be the Poincaré *line bundle* on $J_{\mathbb{Q}} \times J_{\mathbb{Q}}^{\vee}$, trivialised on the union of $\{0\} \times J_{\mathbb{Q}}^{\vee}$ and $J_{\mathbb{Q}} \times \{0\}$. Then the Poincaré *torsor* is the

\mathbb{G}_m -torsor on $J_{\mathbb{Q}} \times J_{\mathbb{Q}}^{\vee}$ defined as

$$(2.1) \quad P_{\mathbb{Q}}^{\times} = \mathbf{Isom}_{J_{\mathbb{Q}} \times J_{\mathbb{Q}}^{\vee}}(\mathcal{O}_{J_{\mathbb{Q}} \times J_{\mathbb{Q}}^{\vee}}, P_{\mathbb{Q}}).$$

For every scheme S over $J_{\mathbb{Q}} \times J_{\mathbb{Q}}^{\vee}$, $P_{\mathbb{Q}}^{\times}(S)$ is the set of isomorphisms from \mathcal{O}_S to $(P_{\mathbb{Q}})_S$, with a free and transitive action of $\mathcal{O}_S(S)^{\times}$. Locally on S for the Zariski topology, $(P_{\mathbb{Q}}^{\times})_S$ is trivial, and $P_{\mathbb{Q}}^{\times}$ is represented by a scheme over $J_{\mathbb{Q}} \times J_{\mathbb{Q}}^{\vee}$.

The theorem of the cube gives $P_{\mathbb{Q}}^{\times}$ the structure of a *biextension* of $J_{\mathbb{Q}}$ and $J_{\mathbb{Q}}^{\vee}$ by \mathbb{G}_m , a notion for the details of which we recommend Section I.2.5 of [22], Grothendieck's Exposés VII and VIII [24], and references therein. This means the following. For S a \mathbb{Q} -scheme, x_1 and x_2 in $J_{\mathbb{Q}}(S)$, and y in $J_{\mathbb{Q}}^{\vee}(S)$, the theorem of the cube gives a canonical isomorphism of \mathcal{O}_S -modules

$$(2.2) \quad (x_1, y)^* P_{\mathbb{Q}} \otimes_{\mathcal{O}_S} (x_2, y)^* P_{\mathbb{Q}} = (x_1 + x_2, y)^* P_{\mathbb{Q}}.$$

This induces a morphism of schemes

$$(2.3) \quad (x_1, y)^* P_{\mathbb{Q}}^{\times} \times_S (x_2, y)^* P_{\mathbb{Q}}^{\times} \longrightarrow (x_1 + x_2, y)^* P_{\mathbb{Q}}^{\times}.$$

as follows. For any S -scheme T , and z_1 in $((x_1, y)^* P_{\mathbb{Q}}^{\times})(T)$ and z_2 in $((x_2, y)^* P_{\mathbb{Q}}^{\times})(T)$, we view z_1 and z_2 as nowhere vanishing sections of the invertible \mathcal{O}_T -modules $(x_1, y)^* P_{\mathbb{Q}}$ and $(x_2, y)^* P_{\mathbb{Q}}$. The tensor product of these two then gives an element of $((x_1 + x_2, y)^* P_{\mathbb{Q}}^{\times})(T)$. This gives $P_{\mathbb{Q}}^{\times} \rightarrow J_{\mathbb{Q}}^{\vee}$ the structure of a commutative group scheme, which is an extension of $J_{\mathbb{Q}}$ by \mathbb{G}_m , over the base $J_{\mathbb{Q}}^{\vee}$. We denote this group law, and the one on $J_{\mathbb{Q}} \times J_{\mathbb{Q}}^{\vee}$, as

$$(2.4) \quad \begin{array}{ccc} (z_1, z_2) & \longmapsto & z_1 +_1 z_2 \\ \downarrow & & \downarrow \\ ((x_1, y), (x_2, y)) & \longmapsto & (x_1, y) +_1 (x_2, y) \quad \longlongequal{\quad} \quad (x_1 + x_2, y). \end{array}$$

In the same way, $P_{\mathbb{Q}}^{\times} \rightarrow J_{\mathbb{Q}}$ has a group law $+_2$ that makes it an extension of $J_{\mathbb{Q}}^{\vee}$ by \mathbb{G}_m over the base $J_{\mathbb{Q}}$. In this way, $P_{\mathbb{Q}}^{\times}$ is both the universal extension of $J_{\mathbb{Q}}$ by \mathbb{G}_m as the universal extension of $J_{\mathbb{Q}}^{\vee}$ by \mathbb{G}_m . The final ingredient of the notion of biextension is that the two partial group laws are compatible in the following sense. For any \mathbb{Q} -scheme S , for x_1 and x_2 in $J_{\mathbb{Q}}(S)$, y_1 and y_2 in $J_{\mathbb{Q}}^{\vee}(S)$, and, for all i and j in $\{1, 2\}$, $z_{i,j}$ in $((x_i, y_j)^* P_{\mathbb{Q}}^{\times})(S)$, we have

$$(2.5) \quad \begin{array}{ccc} (z_{1,1} +_1 z_{2,1}) +_2 (z_{1,2} +_1 z_{2,2}) & \longlongequal{\quad} & (z_{1,1} +_2 z_{1,2}) +_1 (z_{2,1} +_2 z_{2,2}) \\ \downarrow & & \downarrow \\ (x_1 + x_2, y_1) +_2 (x_1 + x_2, y_2) & \longlongequal{\quad} & (x_1, y_1 + y_2) +_1 (x_2, y_1 + y_2) \end{array}$$

with the equality in the upper line taking place in $((x_1 + x_2, y_1 + y_2)^* P_{\mathbb{Q}}^{\times})(S)$.

Now we extend the geometry above over \mathbb{Z} . We denote by J^0 the fibrewise connected component of 0 in J , which is an open subgroup scheme of J , and by Φ the quotient J/J^0 , which is an étale (not necessarily separated) group scheme over \mathbb{Z} , with finite fibres, supported on $\mathbb{Z}/n\mathbb{Z}$. Similarly, we let $J^{\vee 0}$ be the fibrewise connected component of J^{\vee} . Theorem 7.1, in Exposé VIII of [24] gives that $P_{\mathbb{Q}}^{\times}$ extends uniquely to a \mathbb{G}_m -biextension

$$(2.6) \quad P^{\times} \longrightarrow J \times J^{\vee 0}$$

(Grothendieck's pairing on component groups is the obstruction to the existence of such an extension). Note that in this case the existence and the uniqueness follow directly from the requirement of extending the rigidification on $J_{\mathbb{Q}} \times \{0\}$. For details see Section 6.7.

Our base point $b \in C(\mathbb{Z})$ gives an embedding $j_b: C_{\mathbb{Q}} \rightarrow J_{\mathbb{Q}}$, which sends, functorially in \mathbb{Q} -schemes S , an element $c \in C_{\mathbb{Q}}(S)$ to the class of the invertible \mathcal{O}_{C_S} -module $\mathcal{O}_{C_S}(c-b)$. Then j_b extends uniquely to a morphism

$$(2.7) \quad j_b: C^{\text{sm}} \longrightarrow J$$

where C^{sm} is the open subscheme of C consisting of points at which C is smooth over \mathbb{Z} . Note that $C_{\mathbb{Q}}(\mathbb{Q}) = C(\mathbb{Z}) = C^{\text{sm}}(\mathbb{Z})$.

Our next step is to lift j_b , at least on certain opens of C^{sm} , to a morphism to a $\mathbb{G}_m^{\rho-1}$ -torsor over J , where ρ is the rank of the free \mathbb{Z} -module $\text{Hom}(J_{\mathbb{Q}}, J_{\mathbb{Q}}^{\vee})^+$, the \mathbb{Z} -module of self-dual morphisms from $J_{\mathbb{Q}}$ to $J_{\mathbb{Q}}^{\vee}$. This torsor will be the product of pullbacks of P^{\times} via morphisms

$$(2.8) \quad (\text{id}, m \cdot \circ \text{tr}_c \circ f): J \rightarrow J \times J^{\vee 0},$$

with $f: J \rightarrow J^{\vee}$ a morphism of group schemes, $c \in J^{\vee}(\mathbb{Z})$, tr_c the translation by c , m the least common multiple of the exponents of all $\Phi(\overline{\mathbb{F}}_p)$ with p ranging over all primes, and $m \cdot$ the multiplication by m map on J^{\vee} . For such a map $m \cdot \circ \text{tr}_c \circ f$, $j_b: C_{\mathbb{Q}} \rightarrow J_{\mathbb{Q}}$ can be lifted to $(\text{id}, m \cdot \circ \text{tr}_c \circ f)^* P_{\mathbb{Q}}^{\times}$ if and only if $j_b^*(\text{id}, m \cdot \circ \text{tr}_c \circ f)^* P_{\mathbb{Q}}^{\times}$ is trivial. The degree of this \mathbb{G}_m -torsor on $C_{\mathbb{Q}}$ is minus the trace of $\lambda^{-1} \circ m \cdot \circ (f + f^{\vee})$ acting on $H_1(J(\mathbb{C}), \mathbb{Z})$. For example, for $f = \lambda$ the degree is $-4mg$. Note that $j_b: C_{\mathbb{Q}} \rightarrow J_{\mathbb{Q}}$ induces

$$(2.9) \quad j_b^* = -\lambda^{-1}: J_{\mathbb{Q}}^{\vee} \rightarrow J_{\mathbb{Q}},$$

(see [21], Propositions 2.7.9 and 2.7.10). This implies that for f such that this degree is zero, there is a unique c such that $j_b^*(\text{id}, \text{tr}_c \circ f)^* P_{\mathbb{Q}}^{\times}$ is trivial on $C_{\mathbb{Q}}$, and hence also its m th power $j_b^*(\text{id}, m \cdot \circ \text{tr}_c \circ f)^* P_{\mathbb{Q}}^{\times}$.

The map

$$(2.10) \quad \text{Hom}(J_{\mathbb{Q}}, J_{\mathbb{Q}}^{\vee}) \longrightarrow \text{Pic}(J_{\mathbb{Q}}) \longrightarrow \text{NS}_{J_{\mathbb{Q}}/\mathbb{Q}}(\mathbb{Q}) = \text{Hom}(J_{\mathbb{Q}}, J_{\mathbb{Q}}^{\vee})^+$$

sending f to the class of $(\text{id}, f)^* P_{\mathbb{Q}}$ sends f to $f + f^{\vee}$, hence its kernel is $\text{Hom}(J_{\mathbb{Q}}, J_{\mathbb{Q}}^{\vee})^-$, the group of antisymmetric morphisms. But actually, for f antisymmetric, its image in $\text{Pic}(J_{\mathbb{Q}})$ is already zero (see for example [6] and the references therein). Hence the image of $\text{Hom}(J_{\mathbb{Q}}, J_{\mathbb{Q}}^{\vee})$ in $\text{Pic}(J_{\mathbb{Q}})$ is free of rank ρ , and its subgroup of classes with degree zero on $C_{\mathbb{Q}}$ is free of rank $\rho-1$. Let $f_1, \dots, f_{\rho-1}$ be elements of $\text{Hom}(J_{\mathbb{Q}}, J_{\mathbb{Q}}^{\vee})$ whose images in $\text{Pic}(J_{\mathbb{Q}})$ form a basis of this subgroup, and let $c_1, \dots, c_{\rho-1}$ be the corresponding elements of $J^{\vee}(\mathbb{Z})$.

By construction, for each i , the morphism $j_b: C_{\mathbb{Q}} \rightarrow J_{\mathbb{Q}}$ lifts to $(\text{id}, m \cdot \circ \text{tr}_{c_i} \circ f_i)^* P_{\mathbb{Q}}^{\times}$, unique up to \mathbb{Q}^{\times} . Now we spread this out over \mathbb{Z} , to open subschemes U of C^{sm} obtained by removing, for each q dividing n , all but one irreducible components of $C_{\mathbb{F}_q}^{\text{sm}}$, with the remaining irreducible component geometrically irreducible. For such a U , the morphism $\text{Pic}(U) \rightarrow \text{Pic}(C_{\mathbb{Q}})$ is an isomorphism, and $\mathcal{O}_C(U) = \mathbb{Z}$, thus, for each i , there is a lift

$$(2.11) \quad \begin{array}{ccc} & (\text{id}, m \cdot \circ \text{tr}_{c_i} \circ f_i)^* P^{\times} & \\ & \nearrow \tilde{j}_b & \downarrow \\ U & \xrightarrow{j_b} & J \end{array}$$

unique up to $\mathbb{Z}^\times = \{1, -1\}$.

At this point we can explain the strategy of our approach to the quadratic Chabauty method. Let T be the $\mathbb{G}_m^{\rho-1}$ -torsor on J obtained by taking the product of all $T_i := (\text{id}, m \cdot \text{tr}_{c_i} \circ f_i)^* P^\times$:

$$(2.12) \quad \begin{array}{ccc} & T & \longrightarrow P^{\times, \rho-1} \\ & \tilde{j}_b \nearrow & \downarrow \\ U & \xrightarrow{j_b} J & \xrightarrow{(\text{id}, m \cdot \text{tr}_{c_i} \circ f_i)_i} J \times (J^{\vee 0})^{\rho-1} \end{array}$$

Then each $c \in C_{\mathbb{Q}}(\mathbb{Q}) = C^{\text{sm}}(\mathbb{Z})$ lies in one of the finitely many $U(\mathbb{Z})$'s. For each U , we have a lift $\tilde{j}_b: U \rightarrow T$, and, for each prime number p , $\tilde{j}_b(U(\mathbb{Z}))$ is contained in the intersection, in $T(\mathbb{Z}_p)$, of $\tilde{j}_b(U(\mathbb{Z}_p))$ and the closure $\overline{T(\mathbb{Z})}$ of $T(\mathbb{Z})$ in $T(\mathbb{Z}_p)$ with the p -adic topology. Of course, one expects this closure to be of dimension at most $r := \text{rank}(J(\mathbb{Q}))$, and therefore one expects this method to be successful if $r < g + \rho - 1$, the dimension of $T(\mathbb{Z}_p)$. The next two sections make this strategy precise, giving first the necessary p -adic formal and analytic geometry, and then the description of $\overline{T(\mathbb{Z})}$ as a finite disjoint union of images of \mathbb{Z}_p^r under maps constructed from the biextension structure.

3 From algebraic geometry to formal geometry

Let p be a prime number. Given X a smooth scheme of relative dimension d over \mathbb{Z}_p and $x \in X(\mathbb{F}_p)$ let us describe the set $X(\mathbb{Z}_p)_x$ of elements of $X(\mathbb{Z}_p)$ whose image in $X(\mathbb{F}_p)$ is x . The smoothness implies that the maximal ideal of $\mathcal{O}_{X,x}$ is generated by p together with d other elements t_1, \dots, t_d . In this case we call p, t_1, \dots, t_d *parameters at x* ; if moreover $x_l \in X(\mathbb{Z}_p)_x$ is a lift of x such that $t_1(x_l) = \dots = t_d(x_l) = 0$ then we say that the t_i 's are *parameters at x_l* . The t_i can be evaluated on all the points in $X(\mathbb{Z}_p)_x$, inducing a bijection $t := (t_1, \dots, t_d): X(\mathbb{Z}_p)_x \rightarrow (p\mathbb{Z}_p)^d$. We get a bijection

$$(3.1) \quad \tilde{t} := (\tilde{t}_1, \dots, \tilde{t}_d) = \left(\frac{t_1}{p}, \dots, \frac{t_d}{p} \right) : X(\mathbb{Z}_p)_x \xrightarrow{\sim} \mathbb{Z}_p^d.$$

This bijection can be geometrically interpreted as follows. Let $\pi: \tilde{X}_x \rightarrow X$ denote the blow up of X in x . By shrinking X , X is affine and the t_i are regular on X , $t: X \rightarrow \mathbb{A}_{\mathbb{Z}_p}^d$ is etale, and $t^{-1}\{0_{\mathbb{F}_p}\} = \{x\}$. Then $\pi: \tilde{X}_x \rightarrow X$ is the pull back of the blow up of $\mathbb{A}_{\mathbb{Z}_p}^d$ at the origin over \mathbb{F}_p . The affine open part \tilde{X}_x^p of \tilde{X}_x where p generates the image of the ideal m_x of x is the pullback of the corresponding open part of the blow up of $\mathbb{A}_{\mathbb{Z}_p}^d$, which is the multiplication by p morphism $\mathbb{A}_{\mathbb{Z}_p}^d \rightarrow \mathbb{A}_{\mathbb{Z}_p}^d$ that corresponds to $\mathbb{Z}_p[t_1, \dots, t_d] \rightarrow \mathbb{Z}_p[\tilde{t}_1, \dots, \tilde{t}_d]$ with $t_i \mapsto p\tilde{t}_i$. It follows that the p -adic completion $\mathcal{O}(\tilde{X}_x^p)^{\wedge p}$ of $\mathcal{O}(\tilde{X}_x^p)$ is the p -adic completion $\mathbb{Z}_p\langle \tilde{t}_1, \dots, \tilde{t}_d \rangle$ of $\mathbb{Z}_p[\tilde{t}_1, \dots, \tilde{t}_d]$. Explicitly, we have

$$(3.2) \quad \mathbb{Z}_p\langle \tilde{t}_1, \dots, \tilde{t}_d \rangle = \left\{ \sum_{I \in \mathbb{N}^d} a_I \tilde{t}^I \in \mathbb{Z}_p[[\tilde{t}_1, \dots, \tilde{t}_d]] : \forall n \geq 0, \forall^{\text{almost}} I, v_p(a_I) \geq n \right\}.$$

With these definitions, we have

$$(3.3) \quad \begin{aligned} X(\mathbb{Z}_p)_x &= \tilde{X}_x^p(\mathbb{Z}_p) = \text{Hom}(\mathbb{Z}_p\langle \tilde{t}_1, \dots, \tilde{t}_d \rangle, \mathbb{Z}_p) = \mathbb{A}^d(\mathbb{Z}_p), \\ (\tilde{X}_x^p)_{\mathbb{F}_p} &= \text{Spec}(\mathbb{F}_p[\tilde{t}_1, \dots, \tilde{t}_d]). \end{aligned}$$

The affine space $(\tilde{X}_x^p)_{\mathbb{F}_p}$ is canonically a torsor under the tangent space of $X_{\mathbb{F}_p}$ at x .

This construction is functorial. Let Y be a smooth \mathbb{Z}_p -scheme, $f: X \rightarrow Y$ a morphism over \mathbb{Z}_p , and $y := f(x) \in Y(\mathbb{F}_p)$. Then the ideal in $\mathcal{O}_{\tilde{X}_x^p}$ generated by the image of $m_{f(x)}$ is generated by p . That gives us a morphism $\tilde{X}_x^p \rightarrow \tilde{Y}_{f(x)}^p$, and then a morphism from $\mathcal{O}(\tilde{Y}_{f(x)}^p)^{\wedge p}$ to $\mathcal{O}(\tilde{X}_x^p)^{\wedge p}$. Reduction mod p then gives a morphism $(\tilde{X}_x^p)_{\mathbb{F}_p} \rightarrow (\tilde{Y}_{f(x)}^p)_{\mathbb{F}_p}$, the tangent map of f at x , up to a translation.

If this tangent map is injective, and d_x and d_y denote the dimensions of $X_{\mathbb{F}_p}$ at x and of $Y_{\mathbb{F}_p}$ at y , then there are t_1, \dots, t_{d_y} in $\mathcal{O}_{Y,y}$ such that p, t_1, \dots, t_{d_y} are parameters at y , and such that $t_{d_x+1}, \dots, t_{d_y}$ generate the kernel of $\mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{X,x}$. Then the images in $\mathcal{O}_{X,x}$ of p, t_1, \dots, t_{d_x} are parameters at x , and $\mathcal{O}(\tilde{Y}_{f(x)}^p)^{\wedge p} \rightarrow \mathcal{O}(\tilde{X}_x^p)^{\wedge p}$ is $\mathbb{Z}_p\langle \tilde{t}_1, \dots, \tilde{t}_{d_y} \rangle \rightarrow \mathbb{Z}_p\langle \tilde{t}_1, \dots, \tilde{t}_{d_x} \rangle$, with kernel generated by $\tilde{t}_{d_x+1}, \dots, \tilde{t}_{d_y}$.

4 Integral points, closure and finiteness

Let us now return to our original problem. The notation $U, J, T, j_b, \tilde{j}_b, r, \rho$ etc., is as at the end of Section 2. We assume moreover that p does not divide n (n as in the start of Section 2) and that $p > 2$ (for $p = 2$ everything that follows can probably be adapted by working with residue polydiscs modulo 4).

Let u be in $U(\mathbb{F}_p)$, and $t := \tilde{j}_b(u)$. We want a description of the closure $\overline{T(\mathbb{Z})}_t$ of $T(\mathbb{Z})_t$ in $T(\mathbb{Z}_p)_t$. Using the biextension structure of P^\times , we will produce, for each element of $J(\mathbb{Z})_{j_b(u)}$, an element of $T(\mathbb{Z})$ over it. Not all of these points are in $T(\mathbb{Z})_t$, but we will then produce a subset of $T(\mathbb{Z})_t$ whose closure is $\overline{T(\mathbb{Z})}_t$.

If $T(\mathbb{Z})_t$ is empty then $\overline{T(\mathbb{Z})}_t$ is empty, too. So we assume that we have an element $\tilde{t} \in T(\mathbb{Z})_t$ and we define $x_{\tilde{t}} \in J(\mathbb{Z})$ to be the projection of \tilde{t} . Let $f = (f_1, \dots, f_{\rho-1}): J \rightarrow J^{\vee, \rho-1}$, let $c = (c_1, \dots, c_{\rho-1}) \in J^{\vee, \rho-1}(\mathbb{Z})$. We denote by $P^{\times, \rho-1}$ the product over $J \times (J^{\vee 0})^{\rho-1}$ of the $\rho-1$ \mathbb{G}_m -torsors obtained by pullback of P^\times via the projections to $J \times J^{\vee 0}$; it is a biextension of J and $(J^{\vee 0})^{\rho-1}$ by $\mathbb{G}_m^{\rho-1}$, and $T = (\text{id}, m \cdot \circ \text{tr}_c \circ f)^* P^{\times, \rho-1}$. We choose a basis x_1, \dots, x_r of the free \mathbb{Z} -module $J(\mathbb{Z})_0$, the kernel of $J(\mathbb{Z}) \rightarrow J(\mathbb{F}_p)$. For each $i, j \in \{1, \dots, r\}$ we choose $P_{i,j}$, $R_{i,\tilde{t}}$, and $S_{\tilde{t},j}$ in $P^{\times, \rho-1}(\mathbb{Z})$ whose images in $(J \times (J^{\vee 0})^{\rho-1})(\mathbb{Z})$ are $(x_i, f(mx_j))$, $(x_i, (m \cdot \circ \text{tr}_c \circ f)(x_{\tilde{t}}))$ and $(x_{\tilde{t}}, f(mx_j))$:

$$(4.1) \quad \begin{array}{cccc} P_{i,j} & R_{i,\tilde{t}} & S_{\tilde{t},j} & P^{\times, \rho-1} \\ \downarrow & \downarrow & \downarrow & \downarrow \\ (x_i, f(mx_j)) & (x_i, (m \cdot \circ \text{tr}_c \circ f)(x_{\tilde{t}})) & (x_{\tilde{t}}, f(mx_j)) & J \times (J^{\vee 0})^{\rho-1}. \end{array}$$

For each such choice there are $2^{\rho-1}$ possibilities.

For each $n \in \mathbb{Z}^r$ we use the biextension structure on $P^{\times, \rho-1} \rightarrow J \times (J^{\vee 0})^{\rho-1}$ to define the

following points in $P^{\times, \rho-1}(\mathbb{Z})$, with specified images in $(J \times (J^{V0})^{\rho-1})(\mathbb{Z})$:

$$(4.2) \quad \begin{array}{ccc} A_{\tilde{t}}(n) = \sum_{j=1}^r n_j \cdot_2 S_{\tilde{t},j} & & B_{\tilde{t}}(n) = \sum_{i=1}^r n_i \cdot_1 R_{i,\tilde{t}} \\ \downarrow & & \downarrow \\ \left(x_{\tilde{t}}, \sum_{i=1}^r n_i f(mx_i) \right) & & \left(\sum_{i=1}^r n_i x_i, (m \cdot \circ \text{tr}_c \circ f)(x_{\tilde{t}}) \right) \end{array}$$

$$(4.3) \quad \begin{array}{c} C(n) = \sum_{i=1}^r n_i \cdot_1 \left(\sum_{j=1}^r n_j \cdot_2 P_{i,j} \right) \\ \downarrow \\ \left(\sum_{i=1}^r n_i x_i, \sum_{i=1}^r n_i f(mx_i) \right) \end{array}$$

where \sum_1 and \cdot_1 denote iterations of the first partial group law $+_1$ as in (2.4), and analogously for the second group law. We define, for all $n \in \mathbb{Z}^r$,

$$(4.4) \quad D_{\tilde{t}}(n) := (C(n) +_2 B_{\tilde{t}}(n)) +_1 (A_{\tilde{t}}(n) +_2 \tilde{t}) \in P^{\times, \rho-1}(\mathbb{Z}),$$

which is mapped to

$$(4.5) \quad \left(x_{\tilde{t}} + \sum_{i=1}^r n_i x_i, (m \cdot \circ \text{tr}_c \circ f) \left(x_{\tilde{t}} + \sum_{i=1}^r n_i x_i \right) \right) \in (J \times (J^{V0})^{\rho-1})(\mathbb{Z}).$$

Hence $D_{\tilde{t}}(n)$ is in $T(\mathbb{Z})$, and its image in $J(\mathbb{F}_p)$ is $j_b(u)$. We do not know its image in $T(\mathbb{F}_p)$.

We claim that for n in $(p-1)\mathbb{Z}^r$, $D_{\tilde{t}}(n)$ is in $T(\mathbb{Z})_t$. Let n' be in \mathbb{Z}^r and let $n = (p-1)n'$. Then, in the trivial $\mathbb{F}_p^{\times, \rho-1}$ -torsor $P^{\times, \rho-1}(j_b(u), 0)$, on which $+_2$ is the group law, we have:

$$(4.6) \quad A_{\tilde{t}}(n) = (p-1) \cdot_2 A_{\tilde{t}}(n') = 1 \quad \text{in } \mathbb{F}_p^{\times, \rho-1}.$$

Similarly, in $P^{\times, \rho-1}(0, (m \cdot \circ \text{tr}_c \circ f)(j_b(u))) = \mathbb{F}_p^{\times, \rho-1}$, we have $B_{\tilde{t}}(n) = 1$, and, similarly, in $P^{\times, \rho-1}(0, 0) = \mathbb{F}_p^{\times, \rho-1}$, we have $C(n) = 1$. So, with apologies for the mix of additive and multiplicative notations, in $P^{\times, \rho-1}(\mathbb{F}_p)$ we have

$$(4.7) \quad D_{\tilde{t}}(n) = (1 +_2 1) +_1 (1 +_2 t) = t,$$

mapping to the following element in $(J \times J^{V0, \rho-1})(\mathbb{F}_p)$:

$$(4.8) \quad \begin{aligned} & ((0, 0) +_2 ((0, (m \cdot \circ \text{tr}_c \circ f)(j_b(u)))) +_1 ((j_b(u), 0) +_2 (j_b(u), (m \cdot \circ \text{tr}_c \circ f)(j_b(u)))) \\ & = (j_b(u), (m \cdot \circ \text{tr}_c \circ f)(j_b(u))). \end{aligned}$$

We have proved our claim that $D_{\tilde{t}}(n) \in T(\mathbb{Z})_t$.

So we now have the map

$$(4.9) \quad \kappa_{\mathbb{Z}}: \mathbb{Z}^r \rightarrow T(\mathbb{Z})_t, \quad n \mapsto D_{\tilde{t}}((p-1)n).$$

The following theorem will be proved in Section 5.

4.10 Theorem Let x_1, \dots, x_g be in $\mathcal{O}_{J, j_b(u)}$ such that together with p they form a system of parameters of $\mathcal{O}_{J, j_b(u)}$, and let $v_1, \dots, v_{\rho-1}$ be in $\mathcal{O}_{T, t}$ such that $p, x_1, \dots, x_g, v_1, \dots, v_{\rho-1}$ are parameters of $\mathcal{O}_{T, t}$. As in Section 3 these parameters, divided by p , give a bijection

$$(4.10.1) \quad T(\mathbb{Z}_p)_t \longrightarrow \mathbb{Z}_p^{g+\rho-1}.$$

The composition of $\kappa_{\mathbb{Z}}$ with the map (4.10.1) is given by uniquely determined $\kappa_1, \dots, \kappa_{g+\rho-1}$ in $\mathcal{O}(\mathbb{A}_{\mathbb{Z}_p}^r)^{\wedge p} = \mathbb{Z}_p\langle z_1, \dots, z_r \rangle$. The images in $\mathbb{F}_p[z_1, \dots, z_r]$ of $\kappa_1, \dots, \kappa_g$ are of degree at most 1, and the images of $\kappa_{g+1}, \dots, \kappa_{g+\rho-1}$ are of degree at most 2. The map $\kappa_{\mathbb{Z}}$ extends uniquely to the continuous map

$$(4.10.2) \quad \kappa = (\kappa_1, \dots, \kappa_{g+\rho-1}): \mathbb{A}^r(\mathbb{Z}_p) = \mathbb{Z}_p^r \longrightarrow T(\mathbb{Z}_p)_t.$$

and the image of κ is $\overline{T(\mathbb{Z})}_t$.

Now the moment has come to confront $U(\mathbb{Z}_p)_u$ with $\overline{T(\mathbb{Z})}_t$. We have $\tilde{j}_b: U \rightarrow T$, whose tangent map (mod p) at u is injective (here we use that $C_{\mathbb{F}_p}$ is smooth over \mathbb{F}_p). Then, as at the end of Section 3, $\tilde{j}_b: \tilde{U}_u^p \rightarrow \tilde{T}_t^p$ is, after reduction mod p , an affine linear embedding of codimension $g+\rho-2$, $\tilde{j}_b^*: \mathcal{O}(\tilde{T}_t^p)^{\wedge p} \rightarrow \mathcal{O}(\tilde{U}_u^p)^{\wedge p}$ is surjective and its kernel is generated by elements $f_1, \dots, f_{g+\rho-2}$ (we apologise for using the same letter as for the components of $f: J \rightarrow J^{\vee, \rho-1}$), whose images in $\mathbb{F}_p \otimes \mathcal{O}(\tilde{T}_t^p)$ are of degree at most 1, and such that f_1, \dots, f_{g-1} are in $\mathcal{O}(\tilde{J}_{j_b(u)}^p)^{\wedge p}$. The pullbacks $\kappa^* f_i$ are in $\mathbb{Z}_p\langle z_1, \dots, z_r \rangle$; let I be the ideal in $\mathbb{Z}_p\langle z_1, \dots, z_r \rangle$ generated by them, and let

$$(4.11) \quad A := \mathbb{Z}_p\langle z_1, \dots, z_r \rangle / I.$$

Then the elements of \mathbb{Z}_p^r whose image is in $U(\mathbb{Z}_p)_u$ are zeros of I , hence morphisms of rings from A to \mathbb{Z}_p , and hence from the reduced quotient A_{red} to \mathbb{Z}_p .

4.12 Theorem For $i \in \{1, \dots, g+\rho-2\}$, let $\kappa^* \overline{f}_i$ be the image of $\kappa^* f_i$ in $\mathbb{F}_p[z_1, \dots, z_r]$, and let \overline{I} be the ideal of $\mathbb{F}_p[z_1, \dots, z_r]$ generated by them. Then $\kappa^* \overline{f}_1, \dots, \kappa^* \overline{f}_{g-1}$ are of degree at most 1, and $\kappa^* \overline{f}_g, \dots, \kappa^* \overline{f}_{g+\rho-2}$ are of degree at most 2. Assume that $\overline{A} := A/pA = \mathbb{F}_p[z_1, \dots, z_r] / \overline{I}$ is finite. Then \overline{A} is the product of its localisations \overline{A}_m at its finitely many maximal ideals m . The sum of the $\dim_{\mathbb{F}_p} \overline{A}_m$ over the m such that $\overline{A}/m = \mathbb{F}_p$ is an upper bound for the number of elements of \mathbb{Z}_p^r whose image under κ is in $U(\mathbb{Z}_p)_u$, and also an upper bound for the number of elements of $U(\mathbb{Z})$ with image u in $U(\mathbb{F}_p)$.

Proof As every \overline{f}_i is of degree at most 1 in $x_1, \dots, x_g, v_1, \dots, v_{\rho-1}$, every $\kappa^* \overline{f}_i$ is an \mathbb{F}_p -linear combination of $\kappa_1, \dots, \kappa_{g+\rho-1}$, hence of degree at most 2. For $i < g$, \overline{f}_i is a linear combination of x_1, \dots, x_g , and therefore $\kappa^* \overline{f}_i$ is of degree at most 1.

We claim that A is p -adically complete. More generally, let R be a noetherian ring that is J -adically complete for an ideal J , and let I be an ideal in R . The map from R/I to its J -adic completion $(R/I)^\wedge$ is injective ([1, Thm.10.17]). As J -adic completion is exact on finitely generated R -modules ([1, Prop.10.12]), it sends the surjection $R \rightarrow R/I$ to a surjection $R = R^\wedge \rightarrow (R/I)^\wedge$ (see [1, Prop.10.5] for the equality $R = R^\wedge$). It follows that $R/I \rightarrow (R/I)^\wedge$ is surjective.

Now we assume that \overline{A} is finite. As A is p -adically complete, A is the limit of the system of its quotients by powers of p . These quotients are finite: for every $m \in \mathbb{Z}_{\geq 1}$, $A/p^{m+1}A$ is, as abelian group, an extension of A/pA by a quotient of $A/p^m A$. As \mathbb{Z}_p -module, A is generated by any lift of an \mathbb{F}_p -basis of \overline{A} . Hence A is finitely generated as \mathbb{Z}_p -module.

The set of elements of \mathbb{Z}_p^r whose image under κ is in $U(\mathbb{Z}_p)$ is in bijection with the set of \mathbb{Z}_p -algebra morphisms $\text{Hom}(A, \mathbb{Z}_p)$. As A is the product of its localisations A_m at its maximal ideals, $\text{Hom}(A, \mathbb{Z}_p)$ is the disjoint union of the $\text{Hom}(A_m, \mathbb{Z}_p)$. For each m , $\text{Hom}(A_m, \mathbb{Z}_p)$ has at most $\text{rank}_{\mathbb{Z}_p}(A_m)$ elements, and is empty if $\mathbb{F}_p \rightarrow A/m$ is not an isomorphism. This establishes the upper bound for the number of elements of \mathbb{Z}_p^r whose image under κ is in $U(\mathbb{Z}_p)$. By Theorem 4.10, the elements of $U(\mathbb{Z})$ with image u in $U(\mathbb{F}_p)$ are in $\overline{T(\mathbb{Z})}_t$, and therefore of the form $\kappa(x)$ with $x \in \mathbb{Z}_p^r$ such that $\kappa(x)$ is in $U(\mathbb{Z}_p)_u$. This establishes the upper bound for the number of elements of $U(\mathbb{Z})$ with image u in $U(\mathbb{F}_p)$. \square

We include some remarks to explain how Theorem 4.12 can be used, and what we hope that it can do.

4.13 Remark The $\kappa^* \overline{f}_i$ in Theorem 4.12 can be computed from the reduction $\mathbb{F}_p^r \rightarrow T(\mathbb{Z}/p^2\mathbb{Z})$ of $\kappa_{\mathbb{Z}}$ and (to get the \overline{f}_i) from $\tilde{j}_b: U(\mathbb{Z}/p^2\mathbb{Z})_u \rightarrow T(\mathbb{Z}/p^2\mathbb{Z})_t$. For this, one does not need to treat T and J as schemes, one just computes with $\mathbb{Z}/p^2\mathbb{Z}$ -valued points. Now assume that $r \leq g + \rho - 2$. If, for some prime p , the criterion in Theorem 4.12 fails (that is, \overline{A} is not finite), then one can try the next prime. We hope (but also expect) that one quickly finds a prime p such that \overline{A} is finite for every U and for every u in $U(\mathbb{F}_p)$ such that $\tilde{j}_b(u)$ is in the image of $T(\mathbb{Z}) \rightarrow T(\mathbb{F}_p)$. By the way, note that our notation in Theorem 4.12 does not show the dependence on U and u of \tilde{j}_b , $\kappa_{\mathbb{Z}}$, κ and the \overline{f}_i .

4.14 Remark If $r < g + \rho - 2$ then we think that it is likely (when varying p), for dimension reasons, that, for all $u \in U(\mathbb{F}_p)$, the upper bound in Theorem 4.12 for the number of elements of $U(\mathbb{Z})$ with image u in $U(\mathbb{F}_p)$ is sharp.

4.15 Remark Suppose that $r = g + \rho - 2$. Then we expect, for dimension reasons, that it is likely (when varying p) that, for some $u \in U(\mathbb{F}_p)$, the upper bound in Theorem 4.12 for the number of elements of $U(\mathbb{Z})$ with image u in $U(\mathbb{F}_p)$ is not sharp. Then, as in the classical Chabauty method, one must combine the information gotten from several primes, analogous to ‘Mordell-Weil sieving’. Suppose that we are given a subset B of $U(\mathbb{Z})$ that we want to prove to be equal to $U(\mathbb{Z})$. Let B' be the complement in $U(\mathbb{Z})$ of B . For every prime $p > 2$ not dividing n , Theorem 4.12 gives, interpreting \overline{A} as in the end of the proof of Theorem 4.12, a subset O_p of $J(\mathbb{Z})$, that is a union of cosets for the subgroup $p \cdot \ker(J(\mathbb{Z}) \rightarrow J(\mathbb{F}_p))$, that contains $j_b(B')$. Then one hopes that, taking a large enough finite set S of primes, that the intersection of the O_p for p in S is empty.

5 Parametrisation of integral points, and power series

In this section we give a proof of Theorem 4.10. The main tools here are the formal logarithm and formal exponential of a commutative smooth group scheme over a \mathbb{Q} -algebra ([16], Theo-

rem 1): they give us identities as $n \cdot g = \exp(n \cdot \log g)$ that allow us to extend the multiplication to elements n of \mathbb{Z}_p .

The evaluation map from $\mathbb{Z}_p\langle z_1, \dots, z_n \rangle$ to the set of maps $\mathbb{Z}_p^n \rightarrow \mathbb{Z}_p$ is injective (induction on n , non-zero elements of $\mathbb{Z}_p\langle z \rangle$ have only finitely many zeros in \mathbb{Z}_p).

We say that a map $f: \mathbb{Z}_p^n \rightarrow \mathbb{Z}_p^m$ is *given by integral convergent power series* if its coordinate functions are in $\mathbb{Z}_p\langle z_1, \dots, z_n \rangle = \mathcal{O}(\mathbb{A}_{\mathbb{Z}_p}^n)^{\wedge p}$. This property is stable under composition: composition of polynomials over $\mathbb{Z}/p^k\mathbb{Z}$ gives polynomials.

5.1 Logarithm and exponential

Let p be a prime number, and let G be a commutative group scheme, smooth of relative dimension d over a scheme S smooth over \mathbb{Z}_p , with unit section e in $G(S)$. For any s in $S(\mathbb{F}_p)$, $G(\mathbb{Z}_p)_{e(s)}$ is a group fibred over $S(\mathbb{Z}_p)_s$. The fibres have a natural \mathbb{Z}_p -module structure: $G(\mathbb{Z}_p)_{e(s)}$ is the limit of the $G(\mathbb{Z}/p^n\mathbb{Z})_{e(s)}$ ($n \geq 1$), $S(\mathbb{Z}_p)_s$ is the limit of the $S(\mathbb{Z}/p^n\mathbb{Z})_s$, and for each $n \geq 1$, the fibres of $G(\mathbb{Z}/p^n\mathbb{Z})_{e(s)} \rightarrow S(\mathbb{Z}/p^n\mathbb{Z})_s$ are commutative groups annihilated by p^{n-1} . Let $T_{G/S}$ be the relative (geometric) tangent bundle of G over S . Then its pullback $T_{G/S}(e)$ by e is a vector bundle on S of rank d .

5.1.1 Lemma *In this situation, and with n the relative dimension of S over \mathbb{Z}_p , the formal logarithm and exponential of G base changed to $\mathbb{Q} \otimes \mathcal{O}_{S,s}$ converge to maps*

$$\begin{aligned} \log: \tilde{G}_{e(s)}^p(\mathbb{Z}_p) = G(\mathbb{Z}_p)_{e(s)} &\rightarrow (T_{G/S}(e))(\mathbb{Z}_p)_{0(s)} \\ \exp: \tilde{T}_{G/S}(e)_{0(s)}^p(\mathbb{Z}_p) = (T_{G/S}(e))(\mathbb{Z}_p)_{0(s)} &\rightarrow G(\mathbb{Z}_p)_{e(s)}, \end{aligned}$$

that are each others inverse and, after a choice of parameters for $G \rightarrow S$ at $e(s)$ as in (3.1), are given by $n + d$ elements of $\mathcal{O}(\tilde{G}_{e(s)}^p)^{\wedge p}$ and $n + d$ elements of $\mathcal{O}(\tilde{T}_{G/S}(e)_{0(s)}^p)^{\wedge p}$.

For a in \mathbb{Z}_p and g in $G(\mathbb{Z}_p)_{e(s)}$ we have $a \cdot g = \exp(a \cdot \log g)$, and, after a choice of parameters for $G \rightarrow S$ at $e(s)$, this map $\mathbb{Z}_p \times G(\mathbb{Z}_p)_{e(s)} \rightarrow G(\mathbb{Z}_p)_{e(s)}$ is given by $n + d$ elements of $\mathcal{O}(\mathbb{A}_{\mathbb{Z}_p}^1 \times_{\mathbb{Z}_p} \tilde{G}_{e(s)}^p)^{\wedge p}$. The induced morphism $\mathbb{A}_{\mathbb{F}_p}^1 \times (\tilde{G}_{e(s)}^p)_{\mathbb{F}_p} \rightarrow (\tilde{G}_{e(s)}^p)_{\mathbb{F}_p}$, where $(\tilde{G}_{e(s)}^p)_{\mathbb{F}_p}$ is viewed as the product $T_{S_{\mathbb{F}_p}}(s)$ and $T_{G/S}(e(s))$, is a morphism over $T_{S_{\mathbb{F}_p}}(s)$, bilinear in $\mathbb{A}_{\mathbb{F}_p}^1$ and $T_{G/S}(e(s))$.

Proof Let t_1, \dots, t_n be in $\mathcal{O}_{S,s}$ such that p, t_1, \dots, t_n are parameters at s . Then we have a bijection

$$(5.1.2) \quad \tilde{t}: S(\mathbb{Z}_p)_s \rightarrow \mathbb{Z}_p^n, \quad a \mapsto p^{-1} \cdot (t_1(a), \dots, t_n(a)).$$

Similarly, let x_1, \dots, x_d be generators for the ideal $I_{e(s)}$ of e in $\mathcal{O}_{G,e(s)}$. Then p , the t_i and the x_j together are parameters for $\mathcal{O}_{G,e(s)}$, and give the bijection

$$(5.1.3) \quad (t, x)^\sim: G(\mathbb{Z}_p)_{e(s)} \rightarrow \mathbb{Z}_p^{n+d}, \quad b \mapsto p^{-1} \cdot (t_1(b), \dots, x_d(b)).$$

The dx_i form an $\mathcal{O}_{S,s}$ -basis of $\Omega_{G/S}^1(e)_s$, and so give translation invariant differentials ω_i on $G_{\mathcal{O}_{S,s}}$. As G is commutative, for all i , $d\omega_i = 0$ ([16], Proposition 1.3). We also have the dual $\mathcal{O}_{S,s}$ -basis ∂_i of $T_{G/S}(e)$ and the bijection

$$(5.1.4) \quad (t, x)^\sim: (T_{G/S}(e))(\mathbb{Z}_p)_{0(s)} \rightarrow \mathbb{Z}_p^{n+d}, \quad (a, \sum_i v_i \partial_i) \mapsto p^{-1} \cdot (t_1(a), \dots, t_n(a), v_1, \dots, v_d).$$

Then \log is given by elements \log_i in $(\mathbb{Q} \otimes \mathcal{O}_{S,s})[[x_1, \dots, x_d]]$ whose constant term is 0, uniquely determined (Proposition 1.1 in [16]) by the equality

$$(5.1.5) \quad d \log_i = \omega_i, \quad \text{in } \oplus_j \mathcal{O}_{S,s}[[x_1, \dots, x_d]] \cdot dx_j.$$

Hence the formula from calculus, $\log_i(x) - \log_i(0) = \int_0^1 (t \mapsto tx)^* \omega_i$, gives us that, with

$$(5.1.6) \quad \log_i = \sum_{J \neq 0} \log_{i,J} x^J \quad \text{and} \quad \log_{i,J} \in (\mathbb{Q} \otimes \mathcal{O}_{S,s}),$$

we have, for all i and J , with $|J|$ denoting the total degree of x^J ,

$$(5.1.7) \quad |J| \cdot \log_{i,J} \in \mathcal{O}_{S,s}.$$

The claim about convergence and definition of $\log: G(\mathbb{Z}_p)_{e(s)} \rightarrow (T_{G/S}(e))(\mathbb{Z}_p)_{0(s)}$, is now equivalent to having an analytic bijection $\mathbb{Z}_p^{n+d} \rightarrow \mathbb{Z}_p^{n+d}$ given by

$$(5.1.8) \quad \begin{array}{ccc} G(\mathbb{Z}_p)_{e(s)} & \xrightarrow{\quad ? \quad} & (T_{G/S}(e))(\mathbb{Z}_p)_{0(s)} \\ \downarrow (t,x)^\sim & & \downarrow (t,x)^\sim \\ \mathbb{Z}_p^{n+d} & \xrightarrow{\quad ? \quad} & \mathbb{Z}_p^{n+d} \end{array}$$

$$(a, b) \longmapsto \left(a, p^{-1} \cdot \left(\sum_{J \neq 0} \log_{i,J}(\tilde{t}^{-1}(a))(pb)^J \right)_i \right).$$

We have, for each i ,

$$(5.1.9) \quad p^{-1} \cdot \sum_{J \neq 0} \log_{i,J}(\tilde{t}^{-1}(a))(pb)^J = \sum_{J \neq 0} \frac{p^{|J|-1}}{|J|} (|J| \log_{i,J})(\tilde{t}^{-1}(a)) b^J.$$

For each i , this expression is an element of $\mathbb{Z}_p \langle \tilde{t}_1, \dots, \tilde{t}_n, \tilde{x}_1, \dots, \tilde{x}_d \rangle = \mathcal{O}(\tilde{G}_{e(s)}^p)^{\wedge p}$, even when $p = 2$, because for each J , $|J| \log_{i,J}$ is in $\mathcal{O}_{S,s}$, which is contained in $\mathbb{Z}_p \langle \tilde{t}_1, \dots, \tilde{t}_n \rangle$, and the function $\mathbb{Z}_{\geq 1} \rightarrow \mathbb{Q}_p$, $r \mapsto p^{r-1}/r$ has values in \mathbb{Z}_p and converges to 0. The existence and analyticity of \log is now proved (even for $p = 2$). As $p > 2$, the image of (5.1.9) in $\mathbb{F}_p \otimes \mathcal{O}(\tilde{G}_{e(s)}^p)^{\wedge p}$ is \tilde{x}_i , and on the first n coordinates, \log is the identity, so, by applying Hensel modulo powers of p , \log is invertible, and the inverse is also given by $n + d$ elements of $\mathcal{O}(\tilde{T}_{G/S}(e)_{0(s)}^p)^{\wedge p}$.

The function $\mathbb{Z}_p \times G(\mathbb{Z}_p)_{e(s)} \rightarrow G(\mathbb{Z}_p)_{e(s)}$, $(a, g) \mapsto \exp(a \cdot \log g)$ is a composition of maps given by integral convergent power series, hence it is also of that form. \square

5.2 Parametrisation by power series

The assumptions are as in the beginning of Section 4, in particular, $p > 2$. We have a t in $T(\mathbb{F}_p)$, with image $j_b(u)$ in $J(\mathbb{F}_p)$, and a \tilde{t} in $T(\mathbb{Z})$ lifting t . For every Q in $T(\mathbb{Z})$ mapping to $j_b(u)$ in $J(\mathbb{F}_p)$ there are unique $\varepsilon \in \mathbb{Z}^{\times, \rho-1}$ and $n \in \mathbb{Z}^r$ such that $Q = \varepsilon \cdot D_{\tilde{t}}(n)$: the image of Q in $J(\mathbb{Z})$ is in $J(\mathbb{Z})_{j_b(u)}$, hence differs from the image $x_{\tilde{t}}$ in $J(\mathbb{Z})$ of \tilde{t} by an element of $J(\mathbb{Z})_0$ (with here $0 \in J(\mathbb{F}_p)$), $\sum_i n_i x_i$ for a unique $n \in \mathbb{Z}^r$, hence $D_{\tilde{t}}(n)$ and Q are in $T(\mathbb{Z})$ and have the same image in $J(\mathbb{Z})$, and that gives the unique ε . So we have a bijection

$$(5.2.1) \quad \mathbb{Z}^{\times, \rho-1} \times \mathbb{Z}^r \longrightarrow T(\mathbb{Z})_{j_b(u)} = \{Q \in T(\mathbb{Z}) : Q \mapsto j_b(u) \in J(\mathbb{F}_p)\}, \quad (\varepsilon, n) \mapsto \varepsilon \cdot D_{\tilde{t}}(n).$$

But a problem that we are facing is that the map $\mathbb{Z}^r \rightarrow T(\mathbb{F}_p)_{j_b(u)}$ sending n to the image of $D_{\tilde{t}}(n)$ depends on the (unkown) images of the $P_{i,j}$, $R_{i,\tilde{t}}$ and $S_{\tilde{t},j}$ from (4.1) in $P^{\times,\rho-1}(\mathbb{F}_p)$, and so we do not know for which n and ε the point $\varepsilon \cdot D_{\tilde{t}}(n)$ is in $T(\mathbb{Z})_t$. Luckily we have the $\mathbb{Z}_p^{\times,\rho-1}$ -action on $T(\mathbb{Z}_p)$. Using that $\mathbb{Z}_p^\times = \mathbb{F}_p^\times \times (1 + p\mathbb{Z}_p)$ we have $\mathbb{F}_p^{\times,\rho-1}$ acting on $T(\mathbb{Z}_p)_{j_b(u)}$, compatibly with the torsor structure on $T(\mathbb{F}_p)_{j_b(u)}$. So, for every n in \mathbb{Z}^r there is a unique $\xi(n)$ in $\mathbb{F}_p^{\times,\rho-1}$ such that $\xi(n) \cdot D_{\tilde{t}}(n)$ is in $T(\mathbb{Z}_p)_t$. We define

$$(5.2.2) \quad D'(n) := \xi(n) \cdot D_{\tilde{t}}(n).$$

Then for all n in \mathbb{Z}^r ,

$$(5.2.3) \quad \kappa_{\mathbb{Z}}(n) = D_{\tilde{t}}((p-1) \cdot n) = D'((p-1) \cdot n),$$

because $D_{\tilde{t}}((p-1) \cdot n)$ maps to t in $T(\mathbb{F}_p)$. Moreover for every Q in $T(\mathbb{Z})_t$ there is a unique $n \in \mathbb{Z}^r$ and a unique $\varepsilon \in \mathbb{Z}^{\times,\rho-1}$ such that $Q = \varepsilon \cdot D_{\tilde{t}}(n) = \xi(n) \cdot D_{\tilde{t}}(n) = D'(n)$. Hence

$$(5.2.4) \quad T(\mathbb{Z})_t \subset D'(\mathbb{Z}^r).$$

We will now show that, after any choice of parameters of $\mathcal{O}_{T,t}$ as in Theorem 4.10, D' is given by elements $\kappa'_1, \dots, \kappa'_{g+\rho-1}$ of $\mathcal{O}(\mathbb{A}_{\mathbb{Z}_p}^r)^{\wedge p}$, and then $\kappa_{\mathbb{Z}}$ is given by $\kappa_1, \dots, \kappa_{g+\rho-1}$ with, for all $i \in \{1, \dots, g+\rho-1\}$ and all $a \in \mathbb{Z}_p^r$, $\kappa_i(a) = \kappa'_i((p-1)a)$.

We want a formula for $D'(n)$, so we introduce variants of the $P_{i,j}$, $R_{i,\tilde{t}}$, and $S_{\tilde{t},j}$ as follows. The images in $(J \times (J^{v_0})^{\rho-1})(\mathbb{F}_p)$ of these points are of the form $(0, *)$, $(0, *)$, and $(*, 0)$, respectively. Hence the fibers over them of $P^{\times,\rho-1}$ are rigidified, that is, equal to $\mathbb{F}_p^{\times,\rho-1}$. We define their variants $P'_{i,j}$, $R'_{i,\tilde{t}}$, and $S'_{\tilde{t},j}$ in $P^{\times,\rho-1}(\mathbb{Z}_p)$ to be the unique elements in their orbits under $\mathbb{F}_p^{\times,\rho-1}$ whose images in $P^{\times,\rho-1}(\mathbb{F}_p)$ are equal to the element 1 in $\mathbb{F}_p^{\times,\rho-1}$. Replacing, in (4.2) and (4.3), these $P_{i,j}$, $R_{i,\tilde{t}}$, and $S_{\tilde{t},j}$ by $P'_{i,j}$, $R'_{i,\tilde{t}}$, and $S'_{\tilde{t},j}$ gives variants A' , B' and C' , and using these in (4.4) gives a variant $D'_t(n)$ of 5.2.2. Then, for all n in \mathbb{Z}^r , $D'_t(n)$ and $D'(n)$ (as in (5.2.2)) are equal, because both are in $P^{\times,\rho-1}(\mathbb{Z}_p)_t$, and in the same $\mathbb{F}_p^{\times,\rho-1}$ -orbit. Hence we have, for all n in \mathbb{Z}^r :

$$(5.2.5) \quad \begin{aligned} A'(n) &= \sum_{j=1}^r n_j \cdot_2 S'_{\tilde{t},j}, & B'(n) &= \sum_{i=1}^r n_i \cdot_1 R'_{i,\tilde{t}}, \\ C'(n) &= \sum_{i=1}^r n_i \cdot_1 \left(\sum_{j=1}^r n_j \cdot_2 P'_{i,j} \right), \\ D'(n) &= (C'(n) +_2 B'(n)) +_1 (A'(n) +_2 \tilde{t}). \end{aligned}$$

This shows how the map $n \mapsto D'(n)$ is built up from the two partial group laws $+_1$ and $+_2$ on $P^{\times,\rho-1}$, and the iterations \cdot_1 and \cdot_2 . Lemma 5.1.1 gives that the iterations are given by integral convergent power series. The functoriality in Section 3 gives that the maps induced by $+_1$ and $+_2$ on residue polydisks are given by integral convergent power series. Stability under composition then gives that $n \mapsto D'(n)$ is given by elements $\kappa'_1, \dots, \kappa'_{g+\rho-1}$ of $\mathbb{Z}_p \langle z_1, \dots, z_r \rangle$.

We call the κ'_i the coordinate functions of the extension $D': \mathbb{Z}_p^r \rightarrow T(\mathbb{Z}_p)_t = \mathbb{Z}_p^{g+\rho-1}$, and their images $\bar{\kappa}'_1, \dots, \bar{\kappa}'_{g+\rho-1}$ in $\mathbb{F}_p[z_1, \dots, z_r]$ the mod p coordinate functions, viewed as a morphism $\bar{D}'_{\mathbb{F}_p}: \mathbb{A}_{\mathbb{F}_p}^r \rightarrow \mathbb{A}_{\mathbb{F}_p}^{g+\rho-1}$.

The mod p coordinate functions of $A': \mathbb{Z}_p^r \rightarrow P^{\times, \rho-1}(\mathbb{Z}_p) = \mathbb{Z}_p^{\rho g + \rho - 1}$ (after choosing the necessary parameters) are all of degree at most 1. The same holds for B' . We define

$$(5.2.6) \quad C'_2: \mathbb{Z}^r \times \mathbb{Z}^r \longrightarrow P^{\times, \rho-1}(\mathbb{Z}_p), \quad C'_2(n, m) = \sum_{i=1}^r n_i \cdot_1 \left(\sum_{j=1}^r m_j \cdot_2 P'_{i,j} \right).$$

Then the mod p coordinate functions of C'_2 , elements of $\mathbb{F}_p[x_1, \dots, x_r, y_1, \dots, y_r]$, are linear in the x_i , and in the y_j . Hence of degree at most 2, and the same follows for the mod p coordinate functions of C' . However, as the first ρg parameters for $P^{\times, \rho-1}$ come from $J \times J^{\vee \rho-1}$, and the 1st and 2nd partial group laws there act on different factors, the first ρg mod p coordinate functions of C' are in fact linear. As D' is obtained by summing, using the partial group laws, the results of A' , B and C' , we conclude that $\bar{\kappa}'_1, \dots, \bar{\kappa}'_g$ are of degree at most 1, and the remaining $\bar{\kappa}_j$ are of degree at most 2. The same holds then for all $\bar{\kappa}_j$.

5.3 The p -adic closure

We know from (5.2.3) that $\kappa_{\mathbb{Z}}(\mathbb{Z}^r) = D'((p-1)\mathbb{Z}^r)$. From (4.9) we know that $\kappa_{\mathbb{Z}}(\mathbb{Z}^r) \subset T(\mathbb{Z})_t$. From (5.2.4) we know that $T(\mathbb{Z})_t \subset D'(\mathbb{Z}^r)$. So together we have:

$$(5.3.1) \quad D'((p-1)\mathbb{Z}^r) = \kappa_{\mathbb{Z}}(\mathbb{Z}^r) \subset T(\mathbb{Z})_t \subset D'(\mathbb{Z}^r).$$

We have extended D' to a continuous map $\mathbb{Z}_p^r \rightarrow T(\mathbb{Z}_p)_t$. As \mathbb{Z}_p^r is compact, $D'(\mathbb{Z}_p^r)$ is closed in $T(\mathbb{Z}_p)_t$. As \mathbb{Z}^r and $(p-1)\mathbb{Z}^r$ are dense in \mathbb{Z}_p^r , the closures of their images under D' are both equal to $D'(\mathbb{Z}_p^r)$, and equal to $\kappa(\mathbb{Z}_p^r)$. This finishes the proof of Theorem 4.10.

6 Explicit description of the Poincaré torsor

We give an explicit description of the Poincaré torsor on $J \times J^{\vee, 0}$ and its partial group laws, in terms of divisors and invertible \mathcal{O} -modules on C , first over $\mathbb{Z}[1/n]$, where C is smooth, and then over \mathbb{Z} .

6.1 Norms

Let S be a scheme, $f: S' \rightarrow S$ be finite and locally free, say of rank n . Then $\mathcal{O}_{S'} = f_* \mathcal{O}_{S'}$ (we view $\mathcal{O}_{S'}$ as a sheaf on S) is an \mathcal{O}_S -algebra, locally free as \mathcal{O}_S -module of rank n , and $\mathcal{O}_{S'}^\times$ is a subsheaf of groups of $\mathrm{GL}_{\mathcal{O}_S}(\mathcal{O}_{S'})$. Then the norm morphism is the composition

$$(6.1.1) \quad \begin{array}{ccc} & \text{Norm}_{S'/S} & \\ & \curvearrowright & \\ \mathcal{O}_{S'}^\times & \hookrightarrow \mathrm{GL}_{\mathcal{O}_S}(\mathcal{O}_{S'}) & \xrightarrow{\det} \mathcal{O}_S^\times \end{array}$$

For T an $\mathcal{O}_{S'}^\times$ -torsor (triviality locally on S and S' are equivalent, from the equivalence with invertible $\mathcal{O}_{S'}$ -modules), we let $\mathrm{Norm}_{S'/S}(T)$ be the \mathcal{O}_S^\times -torsor

$$(6.1.2) \quad \mathrm{Norm}_{S'/S}(T) := \mathcal{O}_S^\times \otimes_{\mathcal{O}_{S'}^\times} T = (\mathcal{O}_S^\times \times T) / \mathcal{O}_{S'}^\times,$$

with, for every open U of S , and every $u \in \mathcal{O}_{S'}^\times(U)$, u acting as $(v, t) \mapsto (v \cdot \text{Norm}_{S'/S}(u), u^{-1} \cdot t)$. This is functorial in T : a morphism $\varphi: T_1 \rightarrow T_2$ induces an isomorphism $\text{Norm}_{S'/S}(\varphi)$. It is also functorial for cartesian diagrams $(S'_2 \rightarrow S_2) \rightarrow (S'_1 \rightarrow S_1)$.

For $U \subset S$ open, T an $\mathcal{O}_{S'}^\times$ -torsor, and $t \in T(U)$, we have the isomorphism of $\mathcal{O}_{S'}^\times|_U$ -torsors $\mathcal{O}_{S'}^\times|_U \rightarrow T|_U$ sending 1 to t . Functoriality gives $\text{Norm}_{S'/S}(t)$ in $(\text{Norm}_{S'/S}(T))(U)$, also denoted $1 \otimes t$.

The norm functor (6.1.2) is multiplicative:

$$(6.1.3) \quad \text{Norm}_{S'/S}(T_1 \otimes_{\mathcal{O}_{S'}} T_2) = \text{Norm}_{S'/S}(T_1) \otimes_{\mathcal{O}_S} \text{Norm}_{S'/S}(T_2),$$

such that, if $U \subset S$ is open and t_1 and t_2 are in $T_1(U)$ and $T_2(U)$, then

$$(6.1.4) \quad \text{Norm}_{S'/S}(t_1 \otimes t_2) \mapsto \text{Norm}_{S'/S}(t_1) \otimes \text{Norm}_{S'/S}(t_2).$$

Let \mathcal{L} be an invertible $\mathcal{O}_{S'}$ -module; locally on S , it is free of rank 1 as $\mathcal{O}_{S'}$ -module. This gives us the $\mathcal{O}_{S'}^\times$ -torsor (on S) $\text{Isom}_{\mathcal{O}_{S'}}(\mathcal{O}_{S'}, \mathcal{L})$, which gives back \mathcal{L} as $\mathcal{L} = \mathcal{O}_{S'} \otimes_{\mathcal{O}_{S'}^\times} \text{Isom}_{\mathcal{O}_{S'}}(\mathcal{O}_{S'}, \mathcal{L})$. The norm of \mathcal{L} via $f: S' \rightarrow S$ is then defined as

$$(6.1.5) \quad \text{Norm}_{S'/S}(\mathcal{L}) := \mathcal{O}_S \otimes_{\mathcal{O}_S^\times} \text{Norm}_{S'/S}(\text{Isom}_{\mathcal{O}_{S'}}(\mathcal{O}_{S'}, \mathcal{L})).$$

This construction is functorial for isomorphisms of invertible $\mathcal{O}_{S'}$ -modules.

6.2 Norms along finite relative Cartier divisors

This part is inspired by [17], section 1.1. Let S be a scheme, let $f: X \rightarrow S$ be an S -scheme of finite presentation. A finite effective relative Cartier divisor on $f: X \rightarrow S$ is a closed subscheme D of X that is finite and locally free over S , and whose ideal sheaf I_D is locally generated by a non-zero divisor (equivalently, I_D is locally free of rank 1 as \mathcal{O}_X -module). For such a D and an invertible \mathcal{O}_X -module \mathcal{L} , the norm of \mathcal{L} along D is defined, using (6.1.5), as

$$(6.2.1) \quad \text{Norm}_{D/S}(\mathcal{L}) := \text{Norm}_{D/S}(\mathcal{L}|_D).$$

Then $\text{Norm}_{D/S}(\mathcal{L})$ is functorial for cartesian diagrams $(X' \rightarrow S', \mathcal{L}') \rightarrow (X \rightarrow S, \mathcal{L})$.

6.2.2 Lemma *Let $f: X \rightarrow S$ be a morphism of schemes that is of finite presentation. For D a finite effective relative Cartier divisor on f , the norm functor $\text{Norm}_{D/S}$ in (6.2.1) is multiplicative in \mathcal{L} :*

$$(6.2.3) \quad \text{Norm}_{D/S}(\mathcal{L}_1 \otimes \mathcal{L}_2) = \text{Norm}_{D/S}(\mathcal{L}_1) \otimes_{\mathcal{O}_S} \text{Norm}_{D/S}(\mathcal{L}_2),$$

with, for $U \subset S$ open, $V \subset X$ open, containing $f^{-1}U \cap D$ and $l_i \in \mathcal{L}_i(V)$ generating $\mathcal{L}_i|_V$,

$$(6.2.4) \quad \text{Norm}_{D/S}(l_1 \otimes l_2) = \text{Norm}_{D/S}(l_1) \otimes \text{Norm}_{D/S}(l_2).$$

Let D_1 and D_2 be finite effective relative Cartier divisors on f . Then the ideal sheaf $I_{D_1}I_{D_2} \subset \mathcal{O}_X$ is locally free of rank 1, the closed subscheme $D_1 + D_2$ defined by it is a finite effective relative Cartier divisor on f . The norm functor in (6.2.1) is additive in D :

$$(6.2.5) \quad \text{Norm}_{(D_1+D_2)/S}(\mathcal{L}) = \text{Norm}_{D_1/S}(\mathcal{L}) \otimes_{\mathcal{O}_S} \text{Norm}_{D_2/S}(\mathcal{L}),$$

with, for $U \subset S$ open, $V \subset X$ open, containing $f^{-1}U \cap (D_1 + D_2)$ and $l \in \mathcal{L}(V)$ generating $\mathcal{L}|_{D_1+D_2}$,

$$(6.2.6) \quad \text{Norm}_{(D_1+D_2)/S}(l) = \text{Norm}_{D_1/S}(l) \otimes \text{Norm}_{D_2/S}(l).$$

Proof Let D_1 and D_2 be as stated. If $V \subset X$ is open, and f_i generates $I_{D_i}|_V$, then $f_1 f_2$ generates $(I_{D_1} I_{D_2})|_V$, and this element of $\mathcal{O}_X(V)$ is not a zero-divisor because f_1 and f_2 are not. To show that $D_1 + D_2$ is affine over S , we replace S by an affine open of it, and then reduce to the noetherian case, using the assumption that f is of finite presentation. Then, $(D_1 + D_2)_{\text{red}}$ is the image of $D_{1,\text{red}} \amalg D_{2,\text{red}} \rightarrow X$, and therefore is proper. Hence $D_1 + D_2$ is proper over S , and quasi-finite over S , hence finite over S . The short exact sequence

$$(6.2.7) \quad \begin{array}{ccc} I_{D_2}/I_{D_1+D_2} & \hookrightarrow & \mathcal{O}_{D_1+D_2} \twoheadrightarrow \mathcal{O}_{D_2} \\ \parallel & & \\ (I_{D_2})|_{D_1} & & \end{array}$$

shows that $\mathcal{O}_{D_1+D_2}$ is locally free as \mathcal{O}_S -module, of rank the sum of the ranks of the \mathcal{O}_{D_i} . So $D_1 + D_2$ is a finite effective relative Cartier divisor on $X \rightarrow S$.

We prove (6.2.5), by proving the required statement about sheaves of groups. The diagram

$$(6.2.8) \quad \begin{array}{ccccc} & & \text{Norm}_{(D_1+D_2)/S} & & \\ & \searrow & \text{---} & \swarrow & \\ \mathcal{O}_{D_1+D_2}^\times & \longrightarrow & \mathcal{O}_{D_1}^\times \times \mathcal{O}_{D_2}^\times & \xrightarrow{\text{Norm}_{D_1/S} \times \text{Norm}_{D_2/S}} & \mathcal{O}_S^\times \times \mathcal{O}_S^\times & \longrightarrow & \mathcal{O}_S^\times \\ & & & & & & \\ u & \longmapsto & & & & & \text{Norm}_{D_1/S}(u) \text{Norm}_{D_2/S}(u). \end{array}$$

commutes, because multiplication by u on $\mathcal{O}_{D_1+D_2}$ preserves the short exact sequence (6.2.7), multiplying on the sub and quotient by its images in $\mathcal{O}_{D_1}^\times$ and in $\mathcal{O}_{D_2}^\times$; note that the sub is an invertible \mathcal{O}_{D_1} -module. \square

6.3 Explicit description of the Poincaré torsor of a smooth curve

Let g be in $\mathbb{Z}_{\geq 1}$, let S be a scheme, and $\pi: C \rightarrow S$ be a proper smooth curve, with geometrically connected fibres of genus g , with a section $b \in C(S)$. Let $J \rightarrow S$ be its jacobian. On $C \times_S J$ we have $\mathcal{L}^{\text{univ}}$, the universal invertible \mathcal{O} -module of degree zero on C , rigidified at b .

Let $d \geq 0$, and $C^{(d)}$ the d th symmetric power of $C \rightarrow S$ (we note that the quotient $C^d \rightarrow C^{(d)}$ is finite, locally free of rank $d!$, and commutes with base change on S). Then on $C \times_S C^{(d)}$ we have D , the universal effective relative Cartier divisor on C of degree d . Hence, on $C \times_S J \times_S C^{(d)}$ we have their pullbacks D_J and $\mathcal{L}_{C^{(d)}}^{\text{univ}}$, giving us

$$(6.3.1) \quad \mathcal{N}_d := \text{Norm}_{D_J/(J \times_S C^{(d)})}(\mathcal{L}_{C^{(d)}}^{\text{univ}}).$$

This invertible \mathcal{O} -module \mathcal{N}_d on $J \times_S C^{(d)}$, rigidified at the zero-section of J , gives us a morphism of S -schemes $C^{(d)}$ to $\text{Pic}_{J/S}$. The point db (the divisor d times the base point b) in $C^{(d)}(S)$ is mapped to 0, precisely because $\mathcal{L}^{\text{univ}}$ is rigidified at b , and 6.2.5. Hence there is a unique morphism $\square: C^{(d)} \rightarrow J^\vee = \text{Pic}_{J/S}^0$ such that the pullback of the Poincaré bundle P on $J \times J^\vee$ by $(\text{id}, \square): J \times C^{(d)} \rightarrow J \times J^\vee$, with its rigidifications, is the same as \mathcal{N}_d . The following proposition tells us what the morphism \square is, and the next section tells us what the induced isomorphism is between the fibres of \mathcal{N}_d at points of $J \times C^{(d)}$ with the same image in $J \times_S J$.

6.3.2 Proposition *The pullback of P by $(j_b, j_b^{*, -1}): C \times_S J \rightarrow J \times_S J^\vee$ together with its rigidifications at b and 0, is equal to $\mathcal{L}^{\text{univ}}$.*

Let d be in $\mathbb{Z}_{\geq 0}$. The morphism $\square: C^{(d)} \rightarrow J^\vee = \text{Pic}_{J/S}^0$ is the composition of first $\Sigma: C^{(d)} \rightarrow J$, sending, for every S -scheme T and any D in $C^{(d)}(T)$ to the class of $\mathcal{O}_{C_T}(D - db)$ twisted by the pullback from T that makes it rigidified at b , followed by $j_b^{*, -1}: J \rightarrow J^\vee$. Summarised in a diagram, with $\mathcal{M} := (\text{id} \times j_b^{*, -1})^* P$:

$$(6.3.3) \quad \begin{array}{ccccccc} \mathcal{L}^{\text{univ}} & \longleftarrow & P & \longrightarrow & \mathcal{M} & \xrightarrow{\widetilde{\text{id} \times \Sigma}} & \mathcal{N}_d \\ C \times_S J & \xrightarrow{j_b \times j_b^{*, -1}} & J \times_S J^\vee & \xleftarrow{\text{id} \times j_b^{*, -1}} & J \times_S J & \xleftarrow{\text{id} \times \Sigma} & J \times_S C^{(d)}. \end{array}$$

Then \mathcal{M} , with its rigidifications at $\{0\} \times_S J$ and $J \times_S \{0\}$, is symmetric. For $T \rightarrow S$, x in $J(T)$ given by an invertible \mathcal{O} -module \mathcal{L} on C_T rigidified at b , and $y = \Sigma(D)$ in $J(T)$ given by an effective relative divisor D of degree d on C_T we have

$$(6.3.4) \quad P(x, j_b^{*, -1}(y)) = \mathcal{M}(x, y) = \text{Norm}_{D/T}(\mathcal{L}).$$

For c_1 and c_2 in $C(S)$, we have

$$(6.3.5) \quad \mathcal{M}(j_b(c_1), j_b(c_2)) = c_2^*(\mathcal{O}_C(c_1 - b)) \otimes b^*(\mathcal{O}_C(b - c_1)),$$

and, as invertible \mathcal{O} -modules on $C \times_S C$, with Δ the diagonal and $\text{pr}_0: C \times_S C \rightarrow S$ the structure morphism, we have

$$(6.3.6) \quad (j_b \times j_b)^* \mathcal{M} = \mathcal{O}(\Delta) \otimes \text{pr}_1^* \mathcal{O}(-b) \otimes \text{pr}_2^* \mathcal{O}(-b) \otimes \text{pr}_0^* b^* T_{C/S}.$$

For $d > 2g - 2$, $\widetilde{\text{id} \times \Sigma}$ gives \mathcal{N}_d a descent datum along $\text{id} \times \Sigma$ that gives \mathcal{M} on $J \times_S J$. For T an S -scheme, $x \in J(S)$ given by \mathcal{L} on C_T , rigidified at b , D_1 and D_2 in $C^{(d_1)}(S)$ and $C^{(d_2)}(S)$, the isomorphism

$$(6.3.7) \quad \mathcal{M}(x, \Sigma(D_1 + D_2)) = \mathcal{M}(x, \Sigma(D_1)) \otimes \mathcal{M}(x, \Sigma(D_2))$$

corresponds, via $\widetilde{\text{id} \times \Sigma}$, to

$$(6.3.8) \quad \begin{aligned} \mathcal{N}_{d_1+d_2}(x, D_1 + D_2) &= \text{Norm}_{(D_1+D_2)/T}(\mathcal{L}) = \text{Norm}_{D_1/T}(\mathcal{L}) \otimes \text{Norm}_{D_2/T}(\mathcal{L}) \\ &= \mathcal{N}_{d_1}(x, D_1) \otimes \mathcal{N}_{d_2}(x, D_2), \end{aligned}$$

using Lemma 6.2.2.

For T an S -scheme and x_1 and x_2 in $J(T)$ given by \mathcal{O} -modules \mathcal{L}_1 and \mathcal{L}_2 on C_T , rigidified at b , and D in $C^{(d)}(T)$, the isomorphism

$$(6.3.9) \quad \mathcal{M}(x_1 + x_2, \Sigma(D)) = \mathcal{M}(x_1, \Sigma(D)) \otimes \mathcal{M}(x_2, \Sigma(D))$$

corresponds, via $\widetilde{\text{id} \times \Sigma}$, to

$$(6.3.10) \quad \begin{aligned} \mathcal{N}_d(x_1 + x_2, D) &= \text{Norm}_{D/T}(\mathcal{L}_1 \otimes \mathcal{L}_2) = \text{Norm}_{D/T}(\mathcal{L}_1) \otimes \text{Norm}_{D/T}(\mathcal{L}_2) \\ &= \mathcal{N}_d(x_1, D) \otimes \mathcal{N}_d(x_2, D), \end{aligned}$$

using Lemma 6.2.2.

Proof Let T be an S -scheme, and x be in $J(T)$. Then x corresponds to the invertible \mathcal{O} -module $(\text{id} \times x)^* \mathcal{L}^{\text{univ}}$ on C_T , rigidified at b . Let $z := j_b^{*, -1}(x)$ in $J^\vee(T)$. Then $j_b^*(z) = x$, meaning that the pullback of $(\text{id} \times x)^* P$ on J_T rigidified at 0 by j_b equals $(\text{id} \times x)^* \mathcal{L}^{\text{univ}}$ on C_T rigidified at b . Taking $T := J$ and x the tautological point gives the first claim of the proposition.

The symmetry of \mathcal{M} with its rigidifications follows from [21], (2.7.1) and Lemma 2.7.5, and (2.7.7), using 2.9.

Now we prove (6.3.4). So let T and x be as above, and $y = \Sigma(D)$ in $J(T)$ given by a relative divisor D of degree d on C_T . As $C^d \rightarrow C^{(d)}$ is finite and locally free of rank $d!$, we may and do suppose that D is a sum sections, say $D = \sum_{i=1}^d (c_i)$, with $c_i \in C(T)$. Then we have, functorially:

$$(6.3.11) \quad \begin{aligned} P(x, j_b^{*, -1}(y)) &= P(y, j_b^{*, -1}(x)) = P(\Sigma(D), j_b^{*, -1}(x)) \\ &= P\left(\sum_i j_b(c_i), j_b^{*, -1}(x)\right) = \bigotimes_i P(j_b(c_i), j_b^{*, -1}(x)) \\ &= \bigotimes_i \mathcal{L}^{\text{univ}}(c_i, x) = \bigotimes_i \mathcal{L}(c_i) = \text{Norm}_{D/T}(\mathcal{L}). \end{aligned}$$

Identities (6.3.5) and (6.3.6) follow directly from (6.3.4).

Now we prove the claimed compatibility between (6.3.9) and (6.3.10). We do this by considering the case where \mathcal{L} is universal, that is, base changing to J_T and x the universal point. Then, on J_T , we have 2 isomorphisms from $\text{Norm}_{(D_1+D_2)/J_T}(\mathcal{L})$ to $\text{Norm}_{D_1/J_T}(\mathcal{L}) \otimes \text{Norm}_{D_2/J_T}(\mathcal{L})$. These differ by an element of $\mathcal{O}(J_T)^\times = \mathcal{O}(T)^\times$. Hence it suffices to check that this element equals 1 at $0 \in J(T)$. This amounts to checking that the 2 isomorphisms are equal for $\mathcal{L} = \mathcal{O}_{C_T}$ with the standard rigidification at b . Then, both isomorphisms are the multiplication map $\mathcal{O}_T \otimes_{\mathcal{O}_T} \mathcal{O}_T \rightarrow \mathcal{O}_T$.

The compatibility between (6.3.7) and (6.3.8) is proved analogously. \square

6.3.12 Remark From Proposition 6.3.2 one easily deduces, in that situation, for T an S -scheme, x in $J(T)$ given by an invertible \mathcal{O} -module \mathcal{L} on C_T , and D_1 and D_2 effective relative Cartier divisors on C_T , of the same degree, a canonical isomorphism

$$(6.3.13) \quad \mathcal{M}(x, \Sigma(D_1) - \Sigma(D_2)) = \text{Norm}_{D_1/T}(\mathcal{L}) \otimes \text{Norm}_{D_2/T}(\mathcal{L})^{-1},$$

satisfying the analogous compatibilities as in Proposition 6.3.2. No rigidification of \mathcal{L} at b is needed. In fact, for \mathcal{L}_0 an invertible \mathcal{O}_T -module, we have $\text{Norm}_{D_1/T}(\pi^* \mathcal{L}_0) = \mathcal{L}_0^{\otimes d}$, where $\pi: C_T \rightarrow T$ is the structure morphism and d is the degree of D_1 . Hence the right hand side of (6.3.13) is independent of the choice of \mathcal{L} , given x .

6.4 Explicit isomorphism for norms along equivalent divisors

Let g be in $\mathbb{Z}_{\geq 1}$, let S be a scheme, and $p: C \rightarrow S$ be a proper smooth curve, with geometrically connected fibres of genus g , with a section $b \in C(S)$. Let D_1, D_2 be effective relative Cartier divisors of degree d on C , that we also view as elements of $C^{(d)}(S)$. Recall from Proposition 6.3.2 the morphism $\Sigma: C^{(d)} \rightarrow J$. Then $\Sigma(D_1) = \Sigma(D_2)$ if and only if D_1, D_2 are linearly equivalent in the following sense: locally on S , there exists an f in $\mathcal{O}_C(U)^\times$, with $U := C \setminus (D_1 \cup D_2)$, such that $f \cdot: \mathcal{O}_U \rightarrow \mathcal{O}_U$ extends to an isomorphism $f \cdot: \mathcal{O}_C(D_1) \rightarrow \mathcal{O}_C(D_2)$. In this case, we define $\text{div}(f) = D_2 - D_1$. Proposition 6.3.2 gives us, for each invertible \mathcal{O} -module \mathcal{L} of degree 0 on C rigidified at b (viewed as an element of $J(S)$) specific isomorphisms

$$(6.4.1) \quad \begin{aligned} \text{Norm}_{D_1/S}(\mathcal{L}) &= \mathcal{N}_d(\mathcal{L}, D_1) = \mathcal{M}(\mathcal{L}, \Sigma(D_1)) = \mathcal{M}(\mathcal{L}, \Sigma(D_2)) = \mathcal{N}_d(\mathcal{L}, D_2) \\ &= \text{Norm}_{D_2/S}(\mathcal{L}). \end{aligned}$$

Now we describe explicitly this isomorphism $\text{Norm}_{D_1/S}(\mathcal{L}) \rightarrow \text{Norm}_{D_2/S}(\mathcal{L})$. To do so we first describe *an* isomorphism

$$(6.4.2) \quad \varphi_{\mathcal{L}, D_1, D_2}: \text{Norm}_{D_1/S}(\mathcal{L}) \longrightarrow \text{Norm}_{D_2/S}(\mathcal{L})$$

that is functorial for Cartesian diagrams $(C' \rightarrow S', \mathcal{L}', D'_1, D'_2) \rightarrow (C \rightarrow S, \mathcal{L}, D_1, D_2)$ and then we prove that *this* isomorphism is the one in (6.4.1).

We construct $\varphi_{\mathcal{L}, D_1, D_2}$ locally on S and the functoriality of the construction takes care of making it global. So, suppose that f is as above: $f \in \mathcal{O}_C(U)^\times$, and $f \cdot: \mathcal{O}_U \rightarrow \mathcal{O}_U$ extends to an isomorphism $f \cdot: \mathcal{O}_C(D_1) \rightarrow \mathcal{O}_C(D_2)$. Let $n \in \mathbb{Z}$ with $n > 2g - 2 + 2d$. Then $p_*(\mathcal{L}(nb)) \rightarrow p_*\mathcal{L}(nb)|_{D_1+D_2}$ and $p_*(\mathcal{O}_C(nb)) \rightarrow p_*\mathcal{O}_C(nb)|_{D_1+D_2}$ are surjective, and (still localising on S) $p_*(\mathcal{L}(nb))$ and $p_*(\mathcal{O}_C(nb))$ are free \mathcal{O}_S -modules and $\mathcal{L}(nb)|_{D_1+D_2}$ and $\mathcal{O}_C(nb)|_{D_1+D_2}$ are free $\mathcal{O}_{D_1+D_2}$ -modules of rank 1. Then we have l_0 in $(\mathcal{L}(nb))(C)$ and l_1 in $(\mathcal{O}_C(nb))(C)$ restricting to generators on $D_1 + D_2$. Let $D^- := \text{div}(l_1)$ and $D^+ := \text{div}(l_0)$, and let $V := C \setminus (D^+ + D^-)$. Note that V contains $D_1 + D_2$ and that U contains $D^+ + D^-$. Then, on V , $l := l_0/l_1$ is in $\mathcal{L}(V)$, generates $\mathcal{L}|_{D_1+D_2}$, and multiplication by l is an isomorphism $\cdot l: \mathcal{O}_C(D^+ - D^-) \rightarrow \mathcal{L}$, that is, $\text{div}(l) = D^+ - D^-$. Let

$$(6.4.3) \quad f(\text{div}(l)) = f(D^+ - D^-) := \text{Norm}_{D^+/S}(f|_{D^+}) \cdot \text{Norm}_{D^-/S}(f|_{D^-})^{-1} \in \mathcal{O}_S(S)^\times,$$

and let $\varphi_{\mathcal{L}, l, f}$ be the isomorphism, given in terms of generators

$$(6.4.4) \quad \begin{aligned} \varphi_{\mathcal{L}, l, f}: \text{Norm}_{D_1/S}(\mathcal{L}) &\longrightarrow \text{Norm}_{D_2/S}(\mathcal{L}) \\ \text{Norm}_{D_1/S}(l) &\longmapsto f(\text{div}(l))^{-1} \cdot \text{Norm}_{D_2/S}(l). \end{aligned}$$

Now suppose that we made other choices n', l'_0, l'_1 . Then we get $D^{-'}, D^{+'}, V', l'$ and $\varphi_{\mathcal{L}, l', f}$.

Then there is a unique function $g \in \mathcal{O}_C(V \cap V')^\times$ such that $l' = gl$ in $\mathcal{L}(V \cap V')$. Then

$$\begin{aligned}
(6.4.5) \quad \varphi_{\mathcal{L},l',f}(\text{Norm}_{D_1/S}(l)) &= \varphi_{\mathcal{L},l',f}(\text{Norm}_{D_1/S}(g^{-1}l')) \\
&= \varphi_{\mathcal{L},l',f}(g^{-1}(D_1)\text{Norm}_{D_1/S}(l')) \\
&= g^{-1}(D_1) \cdot \varphi_{\mathcal{L},l',f}(\text{Norm}_{D_1/S}(l')) \\
&= g^{-1}(D_1) \cdot f(\text{div}(l'))^{-1} \cdot \text{Norm}_{D_2/S}(l') \\
&= g^{-1}(D_1) \cdot f(\text{div}(gl))^{-1} \cdot \text{Norm}_{D_2/S}(gl) \\
&= g^{-1}(D_1) \cdot f(\text{div}(g) + \text{div}(l))^{-1} \cdot g(D_2) \cdot \text{Norm}_{D_2/S}(l) \\
&= g^{-1}(D_1) \cdot f(\text{div}(g))^{-1} \cdot g(D_2) \cdot f(\text{div}(l))^{-1} \cdot \text{Norm}_{D_2/S}(l) \\
&= g(\text{div}(f)) \cdot f(\text{div}(g))^{-1} \cdot \varphi_{\mathcal{L},l,f}(\text{Norm}_{D_1/S}(l)) \\
&= \varphi_{\mathcal{L},l,f}(\text{Norm}_{D_1/S}(l)),
\end{aligned}$$

where, in the last step, we used Weil reciprocity, in a generality for which we do not know a reference. The truth in this generality is clear from the classical case by reduction to the universal case, in which the base scheme is integral: take a suitable level structure on J , then consider the universal curve with this level structure, and the universal 4-tuple of effective divisors with the necessary conditions. We conclude that $\varphi_{\mathcal{L},l,f} = \varphi_{\mathcal{L},l',f}$.

Now suppose that f' is in $\mathcal{O}_C(U)^\times$ with $\text{div}(f') = \text{div}(f)$. Then there is a unique $u \in \mathcal{O}_S(S)^\times$ such that $f' = u \cdot f$, and since \mathcal{L} has degree 0 on C

$$\begin{aligned}
(6.4.6) \quad \varphi_{\mathcal{L},l,f'}(\text{Norm}_{D_1/S}(l)) &= (u \cdot f)(\text{div}(l))^{-1} \cdot \text{Norm}_{D_2/S}(l) \\
&= u^{-\deg(\text{div}(l))} f(\text{div}(l))^{-1} \cdot \text{Norm}_{D_2/S}(l) \\
&= f(\text{div}(l))^{-1} \cdot \text{Norm}_{D_2/S}(l) = \varphi_{\mathcal{L},l,f}(\text{Norm}_{D_1/S}(l)).
\end{aligned}$$

Hence $\varphi_{\mathcal{L},l,f'} = \varphi_{\mathcal{L},l,f}$. We define

$$(6.4.7) \quad \varphi_{D_1, D_2, \mathcal{L}}: \text{Norm}_{D_1/S}(\mathcal{L}) \longrightarrow \text{Norm}_{D_2/S}(\mathcal{L})$$

as the isomorphism $\varphi_{\mathcal{L},l,f}$ in (6.4.4) for any local choice of f and l .

We now prove that $\varphi_{\mathcal{L},D_1,D_2}$ is the same as the isomorphism in (6.4.1). We do this, as in the proof of Proposition 6.3.2, by considering the case of the universal \mathcal{L} , that is, we base change via $J \rightarrow S$, and then restricting to $0 \in J(S)$. This amounts to checking that the 2 isomorphisms are equal for $\mathcal{L} = \mathcal{O}_C$ with the standard rigidification at b . In this case, $\text{Norm}_{D_i/S}(\mathcal{O}_C) = \mathcal{O}_S$, with $\text{Norm}_{D_i/S}(1) = 1$. Hence $\varphi_{D_1, D_2, \mathcal{O}_C} = \varphi_{\mathcal{O}_C, 1, f}$ is the identity on \mathcal{O}_S (use (6.4.4)). The other isomorphism is the identity on \mathcal{O}_S because of the rigidifications of \mathcal{M} and \mathcal{N}_d on $0 \times J$ and $0 \times C^{(d)}$.

6.5 Symmetry of the Norm for divisors on smooth curves

Let $C \rightarrow S$ be a proper and smooth curve with geometrically connected fibres. For D_1, D_2 effective relative Cartier divisors on C we define an isomorphism

$$(6.5.1) \quad \varphi_{D_1, D_2}: \text{Norm}_{D_1/S}(\mathcal{O}_C(D_2)) \longrightarrow \text{Norm}_{D_2/S}(\mathcal{O}_C(D_1))$$

that is functorial for cartesian diagrams $(C'/S', D'_1, D'_2) \rightarrow (C/S, D_1, D_2)$.

If suffices to define this isomorphism in the universal case, that is, over the scheme that parametrises all D_1 and D_2 . Let d_1 and d_2 be in $\mathbb{Z}_{\geq 0}$, and let $U := C^{(d_1)} \times_S C^{(d_2)}$, and let D_1 and D_2 be the universal divisors on C_U . Then we have the invertible \mathcal{O}_U -modules $\text{Norm}_{D_1/U}(\mathcal{O}_C(D_2))$ and $\text{Norm}_{D_2/U}(\mathcal{O}_C(D_1))$. The image of $D_1 \cap D_2$ in U is closed, let U^0 be its complement. Then, over U^0 , D_1 and D_2 are disjoint, and the restrictions of $\text{Norm}_{D_1/U}(\mathcal{O}_C(D_2))$ and $\text{Norm}_{D_2/U}(\mathcal{O}_C(D_1))$ are generated by $\text{Norm}_{D_1/U}(1)$ and $\text{Norm}_{D_2/U}(1)$, and there is a unique isomorphism $(\varphi_{D_1, D_2})_{U^0}$ that sends $\text{Norm}_{D_1/U}(1)$ to $\text{Norm}_{D_2/U}(1)$.

We claim that this isomorphism extends to an isomorphism over U . To see it, we base change by $U' \rightarrow U$, where $U' = C^{d_1} \times_S C^{d_2}$, then $U' \rightarrow U$ is finite, locally free of rank $d_1! \cdot d_2!$. Then $D_1 = P_1 + \dots + P_{d_1}$ and $D_2 = Q_1 + \dots + Q_{d_2}$ with the P_i and Q_j in $C(U')$. The complement of the inverse image U'^0 in U' of U^0 is the union of the pullbacks $D_{i,j}$ under $\text{pr}_{i,j}: U' \rightarrow C \times_S C$ of the diagonal, that is, the locus where $P_i = Q_j$. Each $D_{i,j}$ is an effective relative Cartier divisor on U' , isomorphic as S -scheme to $C^{d_1+d_2-1}$, hence smooth over S . Now

$$(6.5.2) \quad \text{Norm}_{D_1/U'}(\mathcal{O}(D_2)) = \bigotimes_{i,j} P_i^* \mathcal{O}(Q_j), \quad \text{Norm}_{D_2/U'}(\mathcal{O}(D_1)) = \bigotimes_{i,j} Q_j^* \mathcal{O}(P_i),$$

and, on U'^0 ,

$$(6.5.3) \quad \text{Norm}_{D_1/U'}(1) = \bigotimes_{i,j} 1, \quad \text{Norm}_{D_2/U'}(1) = \bigotimes_{i,j} 1, \quad \text{in } \mathcal{O}(U'^0).$$

The divisor on U' of the tensor-factor 1 at (i, j) , both in $\text{Norm}_{D_1/U'}(1)$ and in $\text{Norm}_{D_2/U'}(1)$, is $D_{i,j}$. Therefore, the isomorphism $(\varphi_{D_1, D_2})_{U^0}$ extends, uniquely, to an isomorphism φ_{D_1, D_2} over U' , which descends uniquely to U .

Our description of φ_{D_1, D_2} allows us to compute it in the trivial case where D_1 and D_2 are disjoint. One should be a bit careful in other cases. For example, when $d_1 = d_2 = 1$ and $P = Q$, we have $P^* \mathcal{O}_C(Q) = P^* \mathcal{O}_C(P)$ is the tangent space of $C \rightarrow S$ at P , and hence also at Q , but $\varphi_{P, Q}$ is multiplication by -1 on that tangent space. The reason for that is that the switch automorphism on $C \times_S C$ induces -1 on the normal bundle of the diagonal.

If b is an S -point on C , because of the symmetry in Proposition 6.3.2, using (6.3.13), for D_1, D_2 relative effective divisors on C of degree d_1, d_2 over S we have the following diagram of isomorphisms defining ψ_{D_1, D_2}

$$(6.5.4) \quad \begin{array}{ccc} \mathcal{M}(\Sigma(D_2), \Sigma(D_1)) & \xlongequal{\quad} & \text{Norm}_{D_1/S}(\mathcal{O}_C(D_2 - d_2b)) \otimes b^* \mathcal{O}_C(D_2 - d_2b)^{-d_1} \\ \parallel & & \downarrow \psi_{D_1, D_2} \\ \mathcal{M}(\Sigma(D_1), \Sigma(D_2)) & \xlongequal{\quad} & \text{Norm}_{D_2/S}(\mathcal{O}_C(D_1 - d_1b)) \otimes b^* \mathcal{O}_C(D_1 - d_1b)^{-d_2}. \end{array}$$

Then

$$(6.5.5) \quad \psi_{D_1, D_2} = \varphi_{D_1, D_2} \otimes \varphi_{D_1, d_2b}^{-1} \otimes \varphi_{d_1b, D_2}^{-1} \otimes \varphi_{d_1b, d_2b}.$$

It is enough to prove it in the universal case, that is when D_1 and D_2 are the universal divisors on C_U , and there we know that there exists a u in $\mathcal{O}_U(U)^\times = \mathcal{O}_S(S)^\times$ such that

$$(6.5.6) \quad u \cdot \psi_{D_1, D_2} = \varphi_{D_1, D_2} \otimes \varphi_{D_1, d_2b}^{-1} \otimes \varphi_{d_1b, D_2}^{-1} \otimes \varphi_{d_1b, d_2b}.$$

Since the symmetry in Proposition 6.3.2 is compatible with the rigidification at $(0, 0) \in (J \times J)(S)$ then $\psi_{d_1 b, d_2 b}$ is the identity on \mathcal{O}_U , as well as the right hand side of (6.5.5) when $D_i = d_i b$. Hence $u = u(d_1 b, d_2 b) = 1$, proving (6.5.5).

Moreover the isomorphisms φ_{D_1, D_2} , and consequently ψ_{D_1, D_2} , are compatible with addition of divisors, that is, under (6.3.10) and (6.3.8), for every triple D_1, D_2, D_3 of relative Cartier divisors on C we have

$$(6.5.7) \quad \varphi_{D_1+D_2, D_3} = \varphi_{D_1, D_3} \otimes \varphi_{D_2, D_3}, \quad \varphi_{D_1, D_2+D_3} = \varphi_{D_1, D_2} \otimes \varphi_{D_1, D_3}.$$

As for (6.5.5), it is enough to prove it in the universal case and then we can reduce to the case where $D_1 = d_1 b$, $D_2 = d_2 b$ and $D_3 = d_3 b$ for d_i positive integers where we have

$$(6.5.8) \quad \begin{aligned} \varphi_{d_1 b + d_2 b, d_3 b} &= \varphi_{d_1 b, d_3 b} \otimes \varphi_{d_2 b, d_3 b} = (-1)^{(d_1 + d_2) d_3}, \\ \varphi_{d_1 b, d_2 b + d_3 b} &= \varphi_{d_1 b, d_2 b} \otimes \varphi_{d_1 b, d_3 b} = (-1)^{d_1 (d_2 + d_3)}. \end{aligned}$$

6.6 Explicit residue disks and partial group laws

Let C be a smooth curve over \mathbb{Z}/p^2 , let $b \in C(\mathbb{Z}/p^2)$, and let \mathcal{M} be as in Proposition 6.3.2. Let $D = D^+ - D^-$ and $E = E^+ - E^-$ be relative Cartier divisors of degree 0 on C . For each α in $\mathcal{M}^\times(\mathbb{F}_p)$ whose image in $(J \times J)(\mathbb{F}_p)$ is given by (D, E) we parametrise $\mathcal{M}^\times(\mathbb{Z}/p^2)_\alpha$, under the assumption that there exists a non-special split reduced divisor of degree g on $C_{\mathbb{F}_p}$.

Let b_1, \dots, b_g in $C(\mathbb{Z}/p^2)$ have distinct images \bar{b}_i in $C(\mathbb{F}_p)$ such that $h^0(C_{\mathbb{F}_p}, \bar{b}_1 + \dots + \bar{b}_g) = 1$, and let b_{g+1}, \dots, b_{2g} in $C(\mathbb{Z}/p^2)$ be such that the \bar{b}_{g+i} are distinct and $h^0(C_{\mathbb{F}_p}, \bar{b}_{g+1} + \dots + \bar{b}_{2g}) = 1$. Then the maps

$$(6.6.1) \quad \begin{aligned} f_1: C^g &\longrightarrow J, & (c_1, \dots, c_g) &\longmapsto [\mathcal{O}_C(c_1 + \dots + c_g - (b_1 + \dots + b_g) + D)] \\ f_2: C^g &\longrightarrow J, & (c_1, \dots, c_g) &\longmapsto [\mathcal{O}_C(c_1 + \dots + c_g - (b_{g+1} + \dots + b_{2g}) + E)] \end{aligned},$$

are étale respectively in the points $(b_1, \dots, b_g) \in C^g(\mathbb{F}_p)$ and $(b_{g+1}, \dots, b_{2g}) \in C^g(\mathbb{F}_p)$ and consequently give bijections $C^g(\mathbb{Z}/p^2)_{(b_1, \dots, b_g)} \rightarrow J(\mathbb{Z}/p^2)_{\bar{D}}$ and $C^g(\mathbb{Z}/p^2)_{(b_{g+1}, \dots, b_{2g})} \rightarrow J(\mathbb{Z}/p^2)_{\bar{E}}$. For each point $c \in C(\mathbb{F}_p)$ we choose $x_{D,c}$ a generator of $\mathcal{O}_C(-D)_c$ and $x_c \in \mathcal{O}_{C_{\mathbb{Z}_p, c}}$ so that p, x_c are parameters in c . For each $i = 1, \dots, 2g$ we choose x_{b_i} so that $x_{b_i}(b_i) = 0$. For each (\mathbb{Z}/p^2) -point $c \in C(\mathbb{Z}/p^2)_{\bar{c}}$ with image \bar{c} in $C(\mathbb{F}_p)$ and for each $\lambda \in \mathbb{F}_p$ let c_λ be the unique point in $C(\mathbb{Z}/p^2)_{\bar{c}}$ with $x_{\bar{c}}(c_\lambda) = \lambda p$. Then the map $\lambda \mapsto c_\lambda$ is a bijection $\mathbb{F}_p \rightarrow C(\mathbb{Z}/p^2)_{\bar{c}}$ hence the maps f_1, f_2 induce bijections

$$(6.6.2) \quad \begin{aligned} \mathbb{F}_p^g &\longrightarrow J(\mathbb{Z}/p^2)_{\bar{D}}, & \lambda &\longmapsto D_\lambda := D + (b_{1, \lambda_1} - b_1) + \dots + (b_{g, \lambda_g} - b_g) \\ \mathbb{F}_p^g &\longrightarrow J(\mathbb{Z}/p^2)_{\bar{E}}, & \mu &\longmapsto E_\mu := E + (b_{g+1, \mu_1} - b_{g+1}) + \dots + (b_{2g, \mu_g} - b_{2g}). \end{aligned}$$

Hence $\mathcal{M}^\times(\mathbb{Z}/p^2)_{\bar{D}, \bar{E}}$ is the union of $\mathcal{M}^\times(D_\lambda, E_\mu)$ as λ and μ vary in \mathbb{F}_p^g and by Proposition 6.3.2 and Remark 6.3.12 we have

$$(6.6.3) \quad \begin{aligned} \mathcal{M}(D_\lambda, E_\mu) &= \text{Norm}_{E^+ / (\mathbb{Z}/p^2)}(\mathcal{O}_C(D_\lambda)) \otimes \text{Norm}_{E^- / (\mathbb{Z}/p^2)}(\mathcal{O}_C(D_\lambda))^{-1} \otimes \\ &\otimes \bigotimes_{i=1}^g (b_{g+i, \mu_i}^* \mathcal{O}_C(D_\lambda) \otimes b_{g+i}^* \mathcal{O}_C(D_\lambda)^{-1}). \end{aligned}$$

For each $i \in \{1, \dots, g\}$, $c \in C(\mathbb{Z}/p^2)$ and $\lambda \in \mathbb{F}_p$ we define $x_i(c, \lambda) := 1$ if $\bar{c} \neq \bar{b}_i$ and $x_i(c, \lambda) := x_{b_i} - \lambda p$ if $\bar{c} = \bar{b}_i$, so that $c^* x_i(c, \lambda)^{-1}$ generates $c^* \mathcal{O}(b_{i, \lambda})$. Then, for each $c \in C(\mathbb{Z}/p^2)$ and each $\lambda \in \mathbb{F}_p^g$,

$$(6.6.4) \quad c^* \left(x_{D, c}^{-1} \cdot \prod_{i=1}^g \frac{x_i(c, 0)}{x_i(c, \lambda_i)} \right) \quad \text{generates } c^* \mathcal{O}_C(D_\lambda).$$

We write $E^\pm = E^{0, \pm} + \dots + E^{g, \pm}$ so that $E^{0, \pm}$ is disjoint from $\{\bar{b}_1, \dots, \bar{b}_g\}$, and $E^{i, \pm}$, restricted to $C_{\mathbb{F}_p}$, is supported on \bar{b}_i . Let $x_{D, E}$ be a generator of $\mathcal{O}_C(-D)$ in a neighborhood of $E^+ \cup E^-$. Then, for each λ in \mathbb{F}_p^g ,

$$(6.6.5) \quad \text{Norm}_{E^{0, \pm}/(\mathbb{Z}/p^2)}(x_{D, E}^{-1}) \otimes \bigotimes_{i=1}^g \text{Norm}_{E^{i, \pm}/(\mathbb{Z}/p^2)} \left(x_{D, E}^{-1} \cdot \frac{x_{b_i}}{x_{b_i} - \lambda_i p} \right)$$

generates $\text{Norm}_{E^\pm/(\mathbb{Z}/p^2)}(\mathcal{O}_C(D_\lambda))$. By (6.6.3), (6.6.4) and (6.6.5) we see that, for λ and μ in \mathbb{F}_p^g ,

$$(6.6.6) \quad \begin{aligned} s_{D, E}(\lambda, \mu) := & \text{Norm}_{E^{0, +}/(\mathbb{Z}/p^2)}(x_{D, E}^{-1}) \otimes \bigotimes_{i=1}^g \text{Norm}_{E^{i, +}/(\mathbb{Z}/p^2)} \left(x_{D, E}^{-1} \cdot \frac{x_{b_i}}{x_{b_i} - \lambda_i p} \right) \otimes \\ & \otimes \text{Norm}_{E^{0, -}/(\mathbb{Z}/p^2)}(x_{D, E}^{-1})^{-1} \otimes \bigotimes_{i=1}^g \text{Norm}_{E^{i, -}/(\mathbb{Z}/p^2)} \left(x_{D, E}^{-1} \cdot \frac{x_{b_i}}{x_{b_i} - \lambda_i p} \right)^{-1} \otimes \\ & \otimes \bigotimes_{i=1}^g \left(b_{g+i, \mu_i}^* \left(x_{D, b_{g+i}}^{-1} \cdot \prod_{j=1}^g \frac{x_j(b_{g+i, \mu_i}, 0)}{x_j(b_{g+i, \mu_i}, \lambda_j)} \right) \otimes b_{g+i}^* \left(x_{D, b_{g+i}}^{-1} \cdot \prod_{j=1}^g \frac{x_j(b_{g+i}, 0)}{x_j(b_{g+i}, \lambda_j)} \right)^{-1} \right) \end{aligned}$$

generates the free rank one \mathbb{Z}/p^2 -module $\mathcal{M}(D_\lambda, E_\mu)$. The fibre $\mathcal{M}^\times(\bar{D}, \bar{E})$ over (\bar{D}, \bar{E}) in $(J \times J)(\mathbb{F}_p)$ is an \mathbb{F}_p^\times -torsor, containing $\overline{s_{D, E}(0, 0)}$ are in bijection with the elements ξ in \mathbb{F}_p^\times and are exactly the points $\xi \cdot s_{D, E}(0, 0)$. Using that $(\mathbb{Z}/p^2)^\times = \mathbb{F}_p^\times \times (1 + p\mathbb{F}_p)$, we parametrise, for each $\xi \in \mathbb{F}_p^\times$, the residue disk of $\xi \cdot \overline{s_{D, E}(0, 0)}$ by the bijection

$$(6.6.7) \quad \mathbb{F}_p^\times \times \mathbb{F}_p^g \times \mathbb{F}_p \longrightarrow \mathcal{M}^\times(\mathbb{Z}/p^2)_{\xi \cdot \overline{s_{D, E}(0, 0)}}, \quad (\lambda, \mu, \tau) \longmapsto (1 + p\tau) \cdot \xi \cdot s_{D, E}(\lambda, \mu).$$

Using this parametrization it is easy to describe the two partial group laws on $\mathcal{M}^\times(\mathbb{Z}/p^2)$ when one of the two points we are summing lies over (\bar{D}, \bar{E}) and the other lies over $(\bar{D}, 0)$ or $(0, \bar{E})$. To compute the group law in $J(\mathbb{Z}/p^2)$ we notice that for each $c \in C(\mathbb{Z}/p^2)$ such that $x_c(c) = 0$ and for each $\lambda, \mu \in \mathbb{F}_p$ we have

$$(6.6.8) \quad \frac{x_c^2}{(x_c - \lambda p)(x_c - \mu p)} = \frac{x_c^2}{x_c^2 - \lambda p x_c - \mu p x_c} = \frac{x_c}{x_c - (\lambda + \mu)p}$$

and since these rational functions generate $\mathcal{O}_C(c_\lambda - c + c_\mu - c)$ and $\mathcal{O}_C(c_{\lambda + \mu} - c)$ in a neighborhood of c , we have the *equality* of relative Cartier divisors on C

$$(6.6.9) \quad (c_\lambda - c) + (c_\mu - c) = c_{\lambda + \mu} - c.$$

Hence, under the definition for $\lambda \in \mathbb{F}_p^g$ of

$$(6.6.10) \quad D_\lambda^0 := (b_{1, \lambda_1} - b_1) + \dots + (b_{g, \lambda_g} - b_g), \quad E_\lambda^0 := (b_{g+1, \lambda_1} - b_{g+1}) + \dots + (b_{2g, \lambda_g} - b_{2g}),$$

we have, for all $\lambda, \mu \in \mathbb{F}_p^g$, that $D_\lambda + D_\mu^0 = D_{\lambda+\mu}$ and $E_\lambda + E_\mu^0 = E_{\lambda+\mu}$. Definition 6.6.6, applied with $(D, 0)$ and $(0, E)$, with $x_{0,E} = 1$ and, for every $c \in C(\mathbb{F}_p)$, with $x_{0,c} = 1$, gives, for all λ, μ in \mathbb{F}_p^g , the elements

$$(6.6.11) \quad s_{D,0}(\lambda, \mu) \in \mathcal{M}^\times(D_\lambda, E_\mu^0), \quad s_{0,E}(\lambda, \mu) \in \mathcal{M}^\times(D_\lambda^0, E_\mu).$$

Then (6.6.8) and (6.6.9), together with the equivalence of (6.3.7) and (6.3.8) and the equivalence of (6.3.9) and (6.3.10) in Proposition 6.3.2, give that for all $\lambda, \lambda_1, \lambda_2, \mu, \mu_1, \mu_2$ in \mathbb{F}_p^g

$$(6.6.12) \quad \begin{aligned} s_{D,0}(\lambda, \mu_1) +_2 s_{D,E}(\lambda, \mu_2) &= s_{D,0}(\lambda, \mu_1) \otimes s_{D,E}(\lambda, \mu_2) = s_{D,E}(\lambda, \mu_1 + \mu_2) \\ s_{0,E}(\lambda_1, \mu) +_1 s_{D,E}(\lambda_2, \mu) &= s_{D,0}(\lambda_1, \mu) \otimes s_{D,E}(\lambda_2, \mu) = s_{D,E}(\lambda_1 + \lambda_2, \mu), \end{aligned}$$

and, consequently, for all $\tau_1, \tau_2 \in \mathbb{F}_p$ and $\xi_1, \xi_2 \in \mathbb{F}_p^\times$, that

$$(6.6.13) \quad \begin{aligned} \xi_1(1+\tau_1 p) \cdot s_{D,0}(\lambda, \mu_1) +_2 \xi_2(1+\tau_2 p) \cdot s_{D,E}(\lambda, \mu_2) &= \xi_1(1+\tau_1 p) \xi_2(1+\tau_2 p) \cdot s_{D,E}(\lambda, \mu_1 + \mu_2) \\ &= \xi_1 \xi_2 (1 + (\tau_1 + \tau_2) p) \cdot s_{D,E}(\lambda, \mu_1 + \mu_2), \\ \xi_1(1+\tau_1 p) \cdot s_{0,E}(\lambda_1, \mu) +_1 \xi_2(1+\tau_2 p) \cdot s_{D,E}(\lambda_2, \mu) &= \xi_1 \xi_2 (1 + (\tau_1 + \tau_2) p) \cdot s_{D,E}(\lambda_1 + \lambda_2, \mu). \end{aligned}$$

Let us now prove that the parametrization (6.6.7) is the inverse of a bijection given by parameters on \mathcal{M}^\times analogously to (3.1). Let \mathcal{Q} be the pullback of \mathcal{M} by $f_1 \times f_2$ with f_1 and f_2 as in (6.6.1). Then the lift $\widetilde{f_1 \times f_2}: \mathcal{Q}^\times \rightarrow \mathcal{M}^\times$ is étale at any point $\beta \in \mathcal{Q}(\mathbb{F}_p)$ lying over $\bar{b} = (b_1, \dots, b_{2g}) \in (C^{2g})(\mathbb{F}_p)$ and induces a bijection between $\mathcal{Q}^\times(\mathbb{Z}/p^2)_{\bar{b}}$ and $\mathcal{M}^\times(\mathbb{Z}/p^2)_{(\bar{D}, \bar{E})}$. In particular we can interpret $s_{D,E}(\lambda, \mu)$ as a section of $\mathcal{Q}(b_{1,\lambda_1}, \dots, b_{2g,\mu_g})$ and we can interpret (6.6.7) as a parametrization of $Q^\times(\mathbb{Z}/p^2)_{\xi_{s_{D,E}(0,0)}}$. It is then enough to prove that (6.6.7) is the inverse of a bijection given by parameters on \mathcal{Q}^\times . It comes from the definition of c_ν for $c \in C(\mathbb{Z}/p^2)$ and $\nu \in \mathbb{F}_p$, that the maps $\lambda_i \mu_i: C^{2g}(\mathbb{Z}/p^2)_{\bar{b}}$ are given by parameters in $\mathcal{O}_{C^{2g}, \bar{b}}$ divided by p . In order to see that also the coordinate $\tau: \mathcal{Q}^\times(\mathbb{Z}/p^2)_{\xi_{s_{D,E}(0)}}$ is given by a parameter divided by p it is enough to prove that there is an open subset $U \subset C^{2g}$ containing \bar{b} and a section s trivializing $\mathcal{Q}|_U$ such that $s_{D,E}(\lambda, \mu) = s(b_{1,\lambda_1}, \dots, b_{2g,\mu_g})$. Remark 6.3.12 and (6.5.1) give that

$$(6.6.14) \quad \begin{aligned} \mathcal{Q} &= \bigotimes_{i,j=1}^g \left((\pi_i, \pi_{g+j})^* \mathcal{O}_{C \times C}(\Delta) \right) \\ &\otimes \bigotimes_{i=1}^g \left(\pi_i^* \mathcal{O}_C(E - (b_{g+1} + \dots + b_{2g})) \otimes \pi_{g+i}^* \mathcal{O}_C(D - (b_1 + \dots + b_g)) \right) \\ &\otimes \text{Norm}_{E/\mathbb{Z}/p^2}(\mathcal{O}_C(D - (b_1 + \dots + b_g))) \otimes \bigotimes_{i=1}^g b_{g+i}^* \mathcal{O}_C(D - (b_1 + \dots + b_g))^{-1} \end{aligned}$$

where $\Delta \subset C \times C$ is the diagonal and π_i is the i -th projection $C^g \times C^g \rightarrow C$. We can prove that there is an open subset $U \subset C^g \times C^g$ containing b and a section s trivializing $\mathcal{Q}|_U$ such that $s_{D,E}(\lambda, \mu) = s(b_{1,\lambda_1}, \dots, b_{2g,\mu_g})$, by trivializing each factor of the above tensor product in a neighborhood of b . Let us see it, for example, for the pieces of the form $(\pi_i, \pi_{g+j})^* \mathcal{O}_{C \times C}(\Delta)$. Let π_1, π_2 be the two projections $C \times C \rightarrow C$ and let us consider the divisor Δ : for each pair of points $c_1, c_2 \in C(\mathbb{F}_p)$ the invertible \mathcal{O} -module $\mathcal{O}_{C \times C}(-\Delta)$ is generated by the section $x_{\Delta, c_1, c_2} := 1$ in a neighborhood of (c_1, c_2) if $c_1 \neq c_2$, while it is generated

by the section $x_{\Delta, c_1, c_2} := \pi_1^* x_{c_1} - \pi_2^* x_{c_2}$ in a neighborhood of (c_1, c_2) if $c_1 = c_2$. If we now take $c_1 = b_i, c_2 = b_{g+j} \in C(\mathbb{F}_p)$ we deduce there is a neighborhood U of (b_i, b_{g+j}) such that $x_{\Delta, b_i, b_{g+j}}^{-1}$ generates $\mathcal{O}_{C \times C}(\Delta)|_U$. For each $\lambda, \mu \in \mathbb{F}_p^g$ the point $(b_{i, \lambda_i}, b_{g+j, \mu_j})$ lies in $U(\mathbb{Z}/p^2)$ and the canonical isomorphism $(b_{i, \lambda_i}, b_{g+j, \mu_j})^* \mathcal{O}_{C \times C}(\Delta) = b_{g+j, \mu_j}^* \mathcal{O}(b_{i, \lambda_i})$ sends the generating section $(b_{i, \lambda_i}, b_{j, \mu_j})^* x_{\Delta, c_1, c_2}^{-1}$ to $b_{j, \mu_j}^* x_i(b_{g+j}, \lambda_i)^{-1}$, which is a factor in (6.6.6). This gives a section $s_{i,j}$ trivializing $\left((\pi_i, \pi_{g+j})^* \mathcal{O}_{C \times C}(\Delta) \right)$ in a neighborhood of b . With similar choices we can find sections trivializing the other factors in (6.6.14) in a neighborhood of b and tensoring all such sections we get a section s such that $s_{D,E}(\lambda, \mu) = s(b_{1, \lambda_1}, \dots, b_{2g, \mu_g})$.

6.7 Extension of the Poincaré biextension over Néron models

Let C over \mathbb{Z} be a curve as in Section 2. Let q be a prime number that divides n . We also write C for $C_{\mathbb{Z}_q}$.

Let J be the Néron model over \mathbb{Z}_q of $\text{Pic}_{C/\mathbb{Q}_q}^0$, and J^0 its fibre-wise connected component of 0. On $(J \times_{\mathbb{Z}_q} J)_{\mathbb{Q}_q}$ we have \mathcal{M} as in Proposition 6.3.2, rigidified at $0 \times J_{\mathbb{Q}_q}$ and $J_{\mathbb{Q}_q} \times 0$. We claim that \mathcal{M} , with these rigidifications, extends uniquely to $J \times_{\mathbb{Z}_q} J^0$. We prove this. First of all, $J \times_{\mathbb{Z}_q} J^0$ is regular, hence Weil divisors and Cartier divisors are the same, and every invertible \mathcal{O} -module on $(J \times_{\mathbb{Z}_q} J^0)_{\mathbb{Q}_q}$ has an extension to an invertible \mathcal{O} -module on $J \times_{\mathbb{Z}_q} J^0$. So let \mathcal{M}' be an extension of \mathcal{M} . Any extension \mathcal{M}'' of \mathcal{M} is then of the form $\mathcal{M}'(D)$, with D a divisor on $J \times_{\mathbb{Z}_q} J^0$ with support in $(J \times_{\mathbb{Z}_q} J^0)_{\mathbb{F}_q}$. Such D are \mathbb{Z} -linear combinations of the irreducible components of the $D_i \times_{\mathbb{F}_q} J_{\mathbb{F}_q}^0$, where the D_i are the irreducible components of $J_{\mathbb{F}_q}$. Now $\mathcal{M}'|_{J \times 0}$ extends $\mathcal{M}|_{J_{\mathbb{Q}_q} \times 0}$, hence the rigidification of $\mathcal{M}|_{J_{\mathbb{Q}_q} \times 0}$ is a rational section of $\mathcal{M}'|_{J \times 0}$ whose divisor is a \mathbb{Z} -linear combination of the D_i . It follows that there is exactly one D as above such that the rigidification of \mathcal{M} extends to a rigidification of $\mathcal{M}'(D)$ on $J \times 0$. That rigidification is compatible with a unique rigidification of $\mathcal{M}'(D)$ on $0 \times J^0$. In what follows, we denote this extension $\mathcal{M}'(D)$ of \mathcal{M} to $J \times_{\mathbb{Z}_q} J^0$ by \mathcal{M} . Let us now prove that the \mathbb{G}_m -torsor \mathcal{M}^\times on $J \times_{\mathbb{Z}_q} J^0$ has a unique biextension structure, extending that over \mathbb{Q}_q . Over $J \times_{\mathbb{Z}_q} J \times_{\mathbb{Z}_q} J^0$ we have the invertible \mathcal{O} -modules whose fibres, at a point (x, y, z) (with values in some \mathbb{Z}_q -scheme) are $\mathcal{M}(x+y, z)$ and $\mathcal{M}(x, z) \otimes \mathcal{M}(y, z)$. The biextension structure over \mathbb{Q}_q gives an isomorphism between these, that differs from an isomorphism over \mathbb{Z}_q by a divisor with support over \mathbb{F}_q . But the compatibility with the rigidification of \mathcal{M} over $J \times_{\mathbb{Z}_q} 0$ proves that this divisor is zero. The other partial group law, and the required properties of them follow in the same way. We have now shown that \mathcal{M}^\times extends the Poincaré biextension.

6.8 Explicit description of the extended Poincaré bundle

Let C over \mathbb{Z} be a curve as in Section 2. Let q be a prime number that divides n . We also write C for $C_{\mathbb{Z}_q}$. By [19], Corollary 9.1.24, C is cohomologically flat over \mathbb{Z}_q , which means that for all \mathbb{Z}_q -algebras A , $\mathcal{O}(C_A) = A$. Another reference for this is [23], (6.1.4), (6.1.6) and (7.2.1).

The relative Picard functor $\text{Pic}_{C/\mathbb{Z}_q}$ sends a \mathbb{Z}_q -scheme T to the set of isomorphism classes of $(\mathcal{L}, \text{rig})$ with \mathcal{L} an invertible \mathcal{O} -module on C_T and rig a rigidification at b . By cohomological flatness, such objects are rigid. But if the action of $\text{Gal}(\overline{\mathbb{F}_q}/\mathbb{F}_q)$ on the set of irreducible

components of $C_{\mathbb{F}_q}$ is non-trivial, then $\text{Pic}_{C/\mathbb{Z}_q}$ is not representable by a \mathbb{Z}_q -scheme, only by an algebraic space over \mathbb{Z}_q (see [23], Proposition 5.5). Therefore, to not be annoyed by such inconveniences, we pass to $S := \text{Spec}(\mathbb{Z}_q^{\text{unr}})$, the maximal unramified extension of \mathbb{Z}_q . Then $\text{Pic}_{C/S}$ is represented by a smooth S -scheme, and on $C \times_S \text{Pic}_{C/S}$ there is a universal pair $(\mathcal{L}^{\text{univ}}, \text{rig})$ ([23], Proposition 5.5, and Section 8.0). We note that $\text{Pic}_{C/S} \rightarrow S$ is separated if and only if $C_{\mathbb{F}_q}$ is irreducible.

Let $\text{Pic}_{C/S}^{[0]}$ be the open part of $\text{Pic}_{C/S}$ where $\mathcal{L}^{\text{univ}}$ is of total degree zero on the fibres of $C \rightarrow S$. It contains the open part $\text{Pic}_{C/S}^0$ where $\mathcal{L}^{\text{univ}}$ has degree zero on all irreducible components of $C_{\mathbb{F}_q}$.

Let E be the closure of the 0-section of $\text{Pic}_{C/S}$, as in [23]. It is contained in $\text{Pic}_{C/S}^{[0]}$. By [23], Proposition 5.2, E is represented by an S -group scheme, étale.

By [23], Theorem 8.1.4, or [8], Theorem 9.5.4, the tautological morphism $\text{Pic}_{C/S}^{[0]} \rightarrow J$ is surjective (for the étale topology) and its kernel is E , and so $J = \text{Pic}_{C/S}^{[0]}/E$. Also, the composition $\text{Pic}_{C/S}^0 \rightarrow \text{Pic}_{C/S}^{[0]} \rightarrow J$ induces an isomorphism $\text{Pic}_{C/S}^0 \rightarrow J^0$.

Let C_i , $i \in I$, be the irreducible components of $C_{\mathbb{F}_q}$. Then, as divisors on C , we have

$$(6.8.1) \quad C_{\mathbb{F}_q} = \sum_{i \in I} m_i C_i.$$

For \mathcal{L} an invertible \mathcal{O} -module on $C_{\mathbb{F}_q}$, its multidegree is defined as

$$(6.8.2) \quad \text{mdeg}(\mathcal{L}): I \rightarrow \mathbb{Z}, \quad i \mapsto \deg_{C_i}(\mathcal{L}|_{C_i}),$$

and its total degree is then

$$(6.8.3) \quad \deg(\mathcal{L}) = \sum_{i \in I} m_i \deg_{C_i}(\mathcal{L}|_{C_i}).$$

The multidegree induces a surjective morphism of groups

$$(6.8.4) \quad \text{mdeg}: \text{Pic}_{C/S}(S) \rightarrow \mathbb{Z}^I.$$

Now let $d \in \mathbb{Z}^I$ be a sufficiently large multidegree so that every invertible \mathcal{O} -module \mathcal{L} on $C_{\mathbb{F}_q}$ with $\text{mdeg}(\mathcal{L}) = d$ satisfies $H^1(C_{\mathbb{F}_q}, \mathcal{L}) = 0$ and has a global section whose divisor is finite. Let \mathcal{L}_0 be an invertible \mathcal{O} -module on C , rigidified at b , with $\text{mdeg}(\mathcal{L}_0) = d$. Then over $C \times_S J^0$ we have the invertible \mathcal{O} -module $\mathcal{L}^{\text{univ}} \otimes \mathcal{L}_0$, and its pushforward \mathcal{E} to J^0 . Then \mathcal{E} is a locally free \mathcal{O} -module on J^0 . Let E be the geometric vector bundle over J^0 corresponding to \mathcal{E} . Then over E , \mathcal{E} has its universal section. Let $U \subset E$ be the open subscheme where the divisor of this universal section is finite over J^0 . The J^0 -group scheme \mathbb{G}_m acts freely on U . We define $V := U/\mathbb{G}_m$. As the \mathbb{G}_m -action preserves the invertible \mathcal{O} -module and its rigidification, the morphism $U \rightarrow J^0$ factors through $U \rightarrow V$ and gives a morphism $\Sigma_{\mathcal{L}_0}: V \rightarrow J^0$. Then on $C \times_S V$ we have the universal effective relative Cartier divisor D^{univ} on $C \times_S V \rightarrow V$ of multidegree d , and $\mathcal{L}^{\text{univ}} \otimes \mathcal{L}_0$ together its rigidification at b is (uniquely) isomorphic to $\mathcal{O}_{C \times_S V}(D^{\text{univ}}) \otimes_{\mathcal{O}_V} b^* \mathcal{O}_{C \times_S V}(-D^{\text{univ}})$ with its tautological rigidification at b , in a diagram:

$$(6.8.5) \quad \mathcal{L}^{\text{univ}} \otimes \mathcal{L}_0 \longlongequal{\quad} \mathcal{O}_{C \times_S V}(D^{\text{univ}}) \otimes_{\mathcal{O}_V} b^* \mathcal{O}_{C \times_S V}(-D^{\text{univ}}).$$

Then $\Sigma_{\mathcal{L}_0}$ sends, for T an S -scheme, a T -point D on C_T to $\mathcal{O}_{C_T}(D) \otimes_{\mathcal{O}_T} b^* \mathcal{O}_{C_T}(-D) \otimes_{\mathcal{O}_C} \mathcal{L}_0^{-1}$ with its rigidification at b . Let s_0 be in $\mathcal{L}_0(C)$ such that its divisor D_0 is finite over S , and let $v_0 \in V(S)$ be the corresponding point.

On $\text{Pic}_{C/S}^{[0]} \times_S V \times_S C$ we have the universal $\mathcal{L}^{\text{univ}}$ from $\text{Pic}_{C/S}^{[0]}$ with rigidification at b , and the universal divisor D^{univ} . Then on $\text{Pic}_{C/S}^{[0]} \times_S V$ we have the invertible \mathcal{O} -module $\mathcal{N}_{q,d}$ whose fibre at a T -point $(\mathcal{L}, \text{rig}, D)$ is $\text{Norm}_{D/T}(\mathcal{L}) \otimes_{\mathcal{O}_T} \text{Norm}_{D_0/T}(\mathcal{L})^{-1}$, canonically trivial on $\text{Pic}_{C/S}^{[0]} \times_S v_0$:

$$(6.8.6) \quad \mathcal{N}_{q,d}: \left(\text{Pic}_{C/S}^{[0]} \times_S V \right) (T) \ni (\mathcal{L}, \text{rig}, D) \longmapsto \text{Norm}_{D/T}(\mathcal{L}) \otimes_{\mathcal{O}_T} \text{Norm}_{D_0/T}(\mathcal{L})^{-1}.$$

Any global regular function on the integral scheme $\text{Pic}_{C/S}^{[0]} \times_S V$ is constant on the generic fibre, hence in $\mathbb{Q}_q^{\text{unr}}$, and restricting it to $(0, v_0)$ shows that it is in $\mathbb{Z}_q^{\text{unr}}$, and if it is 1 on $\text{Pic}_{C/S}^{[0]} \times_S v_0$, it is equal to 1. Therefore trivialisations on $\text{Pic}_{C/S}^{[0]} \times_S v_0$ rigidify invertible \mathcal{O} -modules on $\text{Pic}_{C/S}^{[0]} \times_S V$.

The next proposition generalises [21], Corollary 2.8.6 and Lemma 2.7.11.2: there, $C \rightarrow S$ is nodal (but not necessarily regular), and the restriction of \mathcal{M} to $J^0 \times_S J^0$ is described.

6.8.7 Proposition *In the situation of Section 6.8, the pullback of the invertible \mathcal{O} -module \mathcal{M} on $J \times_{\mathbb{Z}_q^{\text{unr}}} J^0$ to $\text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]} \times_{\mathbb{Z}_q^{\text{unr}}} V$ by the product of the quotient map $\text{quot}: \text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]} \rightarrow J$ and the map $\Sigma_{\mathcal{L}_0}: V \rightarrow J^0$ is $\mathcal{N}_{q,d}$, compatible with their rigidifications at $J \times 0$ and $\text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]} \times v_0$. In a diagram:*

$$(6.8.8) \quad \begin{array}{ccccc} P^\times & \longleftarrow & \mathcal{M}^\times & \longleftarrow & \mathcal{N}_{q,d}^\times \\ \downarrow & & \downarrow & & \downarrow \\ J \times_{\mathbb{Z}_q^{\text{unr}}} J^{\vee,0} & \xleftarrow{\text{id} \times j_b^{*, -1}} & J \times_{\mathbb{Z}_q^{\text{unr}}} J^0 & \xleftarrow{\text{quot} \times \Sigma_{\mathcal{L}_0}} & \text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]} \times_{\mathbb{Z}_q^{\text{unr}}} V. \end{array}$$

For T any $\mathbb{Z}_q^{\text{unr}}$ -scheme, for x in $J(T)$ given by an invertible \mathcal{O} -module \mathcal{L} on C_T rigidified at b , and y in $J^0(T) = \text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^0(T)$ given by the difference $D = D^+ - D^-$ of effective relative Cartier divisors on C_T of the same multidegree, we have

$$P(x, j_b^{*, -1}(y)) = \mathcal{M}(x, y) = \text{Norm}_{D^+/T}(\mathcal{L}) \otimes_{\mathcal{O}_T} \text{Norm}_{D^-/T}(\mathcal{L})^{-1}.$$

Proof The scheme $\text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]} \times_{\mathbb{Z}_q^{\text{unr}}} V$ is smooth over $\mathbb{Z}_q^{\text{unr}}$, hence regular, it is connected, hence integral, and since $V_{\mathbb{F}_q}$ is irreducible, the irreducible components of $(\text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]} \times_{\mathbb{Z}_q^{\text{unr}}} V)_{\mathbb{F}_q}$ are the $P^i \times_{\mathbb{F}_q} V_{\mathbb{F}_q}$, with P^i the irreducible components of $(\text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]})_{\mathbb{F}_q}$, with i in $\pi_0((\text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]})_{\mathbb{F}_q})$, which, by the way, equals the kernel of $\mathbb{Z}^I \rightarrow \mathbb{Z}$, $x \mapsto \sum_{j \in I} m_j x_j$.

We prove the first claim. Both $\mathcal{N}_{q,d}$ and the pullback of \mathcal{M} are rigidified on $\text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]} \times v_0$. Below we will give, after inverting q , an isomorphism α from $\mathcal{N}_{q,d}$ to the pullback of \mathcal{M} that is compatible with the rigidifications. Then there is a unique divisor D_α on $\text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]} \times_{\mathbb{Z}_q^{\text{unr}}} V$, supported on $(\text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]} \times_{\mathbb{Z}_q^{\text{unr}}} V)_{\mathbb{F}_q}$, such that α is an isomorphism from $\mathcal{N}_{q,d}(D_\alpha)$ to the pullback of \mathcal{M} . Let i be in $\pi_0((\text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]})_{\mathbb{F}_q})$, and let x be in $\text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]}(\mathbb{Z}_q^{\text{unr}})$ specialising to an \mathbb{F}_q -point

of P^i , then restricting α to (x_i, v_0) and using the compatibility of α (over $\mathbb{Q}_q^{\text{unr}}$) with the rigidifications, gives that the multiplicity of $P^i \times V_{\mathbb{F}_q}$ in D_α is zero. Hence D_α is zero.

Let us now give, over $(\text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]} \times_{\mathbb{Z}_q^{\text{unr}}} V)_{\mathbb{Q}_q^{\text{unr}}}$, an isomorphism α from $\mathcal{N}_{q,d}$ to the pullback of \mathcal{M} . Note that $(\text{Pic}_{C/\mathbb{Z}_q^{\text{unr}}}^{[0]})_{\mathbb{Q}_q^{\text{unr}}} = J_{\mathbb{Q}_q^{\text{unr}}}$, and that $V_{\mathbb{Q}_q^{\text{unr}}} = C_{\mathbb{Q}_q^{\text{unr}}}^{(|d|)}$, where $|d| = \sum_i m_i d_i$ is the total degree given by the multidgree d . For T a $\mathbb{Q}_q^{\text{unr}}$ -scheme, $x \in J(T)$ given by \mathcal{L} an invertible \mathcal{O}_{C_T} -module rigidified at b , and $v \in V(T)$ given by a relative Cartier divisor D of degree $|d|$ on C_T , we have, using Proposition 6.3.2 and (6.8.6), the following isomorphisms (functorial in T), respecting the rigidifications at $v = v_0$:

$$(6.8.9) \quad \begin{aligned} \mathcal{M}(x, \Sigma_{\mathcal{L}_0}(v)) &= \mathcal{M}(x, \Sigma(v) - \Sigma(v_0)) = \mathcal{M}(x, \Sigma(v)) \otimes \mathcal{M}(x, \Sigma(v_0))^{-1} \\ &= \text{Norm}_{D/T}(\mathcal{L}) \otimes_{\mathcal{O}_T} \text{Norm}_{D_0/T}(\mathcal{L})^{-1} = \mathcal{N}_{q,d}(x, v). \end{aligned}$$

This finishes the proof of the first claim of the Proposition. The second claim follows directly from the definition of $\mathcal{N}_{q,d}$, plus the compatibility at the end of Proposition 6.3.2. \square

6.9 Integral points of the extended Poincaré torsor

Let C over \mathbb{Z} be a curve as in Section 2. Given a point $(x, y) \in (J \times J^0)(\mathbb{Z})$ we want to describe explicitly the free \mathbb{Z} -module $\mathcal{M}(x, y)$ when x is given by an invertible \mathcal{O} -module \mathcal{L} of total degree 0 on C rigidified at b and y is given as a relative Cartier divisor D on C of total degree 0 with the property that there exists a unique divisor V whose support is disjoint from b and contained in the bad fibres of $C \rightarrow \text{Spec}(\mathbb{Z})$ such that $\mathcal{O}(D+V)$ has degree zero when restricted to every irreducible component of any fibre of $C \rightarrow \text{Spec}(\mathbb{Z})$. Since $\mathcal{M}(x, y)$ is a free \mathbb{Z} -module of rank 1 then it is a submodule of $\mathcal{M}(x, y)[1/n]$ and writing $D = D^+ - D^-$ as a difference of relative effective Cartier divisors, Proposition 6.3.2, with $S = \text{Spec}(\mathbb{Z}[1/n])$, gives

$$(6.9.1) \quad \mathcal{M}(x, y)[1/n] = (\text{Norm}_{D^+/\mathbb{Z}}(\mathcal{L}) \otimes_{\mathbb{Z}} \text{Norm}_{D^-/\mathbb{Z}}(\mathcal{L})^{-1}) [1/n]$$

and consequently there exist unique integers e_q , for q varying among the primes dividing n , such that, as submodules of $(\text{Norm}_{D^+/\mathbb{Z}}(\mathcal{L}) \otimes_{\mathbb{Z}} \text{Norm}_{D^-/\mathbb{Z}}(\mathcal{L})^{-1}) [1/n]$,

$$(6.9.2) \quad \mathcal{M}(x, y) = \left(\prod_{q|n} q^{e_q} \right) \cdot (\text{Norm}_{D^+/\mathbb{Z}}(\mathcal{L}) \otimes_{\mathbb{Z}} \text{Norm}_{D^-/\mathbb{Z}}(\mathcal{L})^{-1}).$$

We write $V = \sum_{q|n} V_q$ where V_q is a divisor supported on $C_{\mathbb{F}_q}$. For every prime q dividing n let $C_{i,q}$, $i \in I_q$ the irreducible components of $C_{\mathbb{F}_q}$ with multiplicity $m_{i,q}$ and let $V_{i,q}$ be the integers so that $V_q = \sum_{i \in I_q} V_{i,q} C_{i,q}$.

For every q dividing n let H_q be an effective relative Cartier divisor on $C_{\mathbb{Z}_q}$ whose complement U_q is affine (recall that C is projective over \mathbb{Z} , take a high degree embedding and a hyperplane section that avoids chosen closed points $c_{i,q}$ on the $C_{i,q}$). The Chinese remainder theorem, applied to the $\mathcal{O}_C(U_q)$ -module $(\mathcal{O}_C(D+V))(U_q)$ and the (distinct) closed points $c_{i,q}$, provides an element f_q of $(\mathcal{O}_C(D+V))(U_q)$ that generates $\mathcal{O}_C(D+V)$ at all $c_{i,q}$. Let $D_q = D_q^+ - D_q^-$ be the divisor of f_q as rational section of $\mathcal{O}_C(D+V)$. Then D_q^+ and D_q^- are finite over \mathbb{Z}_q , and f_q is a rational function on $C_{\mathbb{Z}_q}$ with

$$(6.9.3) \quad \text{div}(f_q) = (D_q^+ - D_q^-) - (D+V) = (D_q^+ + D^-) - (D^+ + D_q^-) - V.$$

This linear equivalence, restricted to \mathbb{Q}_q , gives the isomorphism (6.4.7)

$$(6.9.4) \quad \varphi: \text{Norm}_{(D^++D_q^-)/\mathbb{Q}_q}(\mathcal{L}) \longrightarrow \text{Norm}_{(D_q^++D^-)/\mathbb{Q}_q}(\mathcal{L}).$$

Tensoring with $\text{Norm}_{(D^-+D_q^-)/\mathbb{Q}_q}(\mathcal{L})^{-1}$ we obtain the isomorphism

$$(6.9.5) \quad \varphi \otimes \text{id}: \text{Norm}_{D^+/\mathbb{Q}_q}(\mathcal{L}) \otimes \text{Norm}_{D^-/\mathbb{Q}_q}(\mathcal{L})^{-1} \longrightarrow \text{Norm}_{D_q^+/\mathbb{Q}_q}(\mathcal{L}) \otimes \text{Norm}_{D_q^-/\mathbb{Q}_q}(\mathcal{L})^{-1}$$

using the identifications

$$(6.9.6) \quad \begin{aligned} \text{Norm}_{D^+/\mathbb{Q}_q}(\mathcal{L}) \otimes \text{Norm}_{D^-/\mathbb{Q}_q}(\mathcal{L})^{-1} &= \text{Norm}_{(D^++D_q^-)/\mathbb{Q}_q}(\mathcal{L}) \otimes \text{Norm}_{(D^-+D_q^-)/\mathbb{Q}_q}(\mathcal{L})^{-1} \\ \text{Norm}_{D_q^+/\mathbb{Q}_q}(\mathcal{L}) \otimes \text{Norm}_{D_q^-/\mathbb{Q}_q}(\mathcal{L})^{-1} &= \text{Norm}_{(D_q^++D^-)/\mathbb{Q}_q}(\mathcal{L}) \otimes \text{Norm}_{(D^-+D_q^-)/\mathbb{Q}_q}(\mathcal{L})^{-1}. \end{aligned}$$

Using the same method as for getting the rational section f_q of $\mathcal{O}_C(D+V)$, we get a rational section l of \mathcal{L} with the support of $\text{div}(l)$ finite over \mathbb{Z}_q and disjoint from the supports of D and D_q , and from the intersections of different $C_{i,q}$ and $C_{j,q}$. By Proposition 6.8.7, and the choice of l ,

$$(6.9.7) \quad \mathcal{M}(x, y)_{\mathbb{Z}_q} = \text{Norm}_{D_q^+/\mathbb{Z}_q}(\mathcal{L}) \otimes \text{Norm}_{D_q^-/\mathbb{Z}_q}(\mathcal{L})^{-1} = \mathbb{Z}_q \cdot \text{Norm}_{D_q^+/\mathbb{Z}_q}(l) \otimes \text{Norm}_{D_q^-/\mathbb{Z}_q}(l)^{-1},$$

and

$$(6.9.8) \quad \text{Norm}_{D^+/\mathbb{Z}_q}(\mathcal{L}) \otimes \text{Norm}_{D^-/\mathbb{Z}_q}(\mathcal{L})^{-1} = \mathbb{Z}_q \cdot \text{Norm}_{D^+/\mathbb{Z}_q}(l) \otimes \text{Norm}_{D^-/\mathbb{Z}_q}(l)^{-1}.$$

By (6.4.4), we have

$$(6.9.9) \quad \varphi \otimes \text{id}: \text{Norm}_{D^+/\mathbb{Q}_q}(l) \otimes \text{Norm}_{D^-/\mathbb{Q}_q}(l)^{-1} \mapsto f_q(\text{div}(l))^{-1} \cdot \text{Norm}_{D_q^+/\mathbb{Q}_q}(l) \otimes \text{Norm}_{D_q^-/\mathbb{Q}_q}(l)^{-1}.$$

Comparing with (6.9.2), we conclude that

$$(6.9.10) \quad e_q = v_q(f_q(\text{div}(l))).$$

We write $\text{div}(l) = \sum_j n_j D_j$ as a sum of prime divisors. These D_j are finite over \mathbb{Z}_q , disjoint from the support of the horizontal part of $\text{div}(f_q)$, that is of $D_q - D$, and each of them meets only one of the $C_{i,q}$, say $C_{s(j),q}$. Then, for each j , $f_q^{m_{s(j),q}}$ and $q^{-V_{s(j),q}}$ have the same multiplicity along $C_{s(j),q}$, and consequently they differ multiplicatively by a unit on a neighborhood of D_j . Then we have

$$(6.9.11) \quad \begin{aligned} v_q(f_q(D_j)) &= \frac{v_q(f_q^{m_{s(j),q}}(D_j))}{m_{s(j),q}} = \frac{v_q(q^{-V_{s(j),q}}(D_j))}{m_{s(j),q}} = \frac{v_q(\text{Norm}_{D_j/\mathbb{Z}_q}(q^{-V_{s(j),q}}))}{m_{s(j),q}} \\ &= \frac{-V_{s(j),q} \deg_{\mathbb{Z}_q}(D_j)}{m_{s(j),q}} = \frac{-V_{s(j),q} \cdot (D_j \cdot C_{s(j),q})}{m_{s(j),q}} = \frac{-V_{s(j),q} \cdot (D_j \cdot m_{s(j),q} C_{s(j),q})}{m_{s(j),q}} \\ &= -V_{s(j),q}(D_j \cdot C_{s(j)}) = -V_q \cdot D_j. \end{aligned}$$

We get

$$(6.9.12) \quad e_q = v_q(f_q(\text{div}(l))) = -V_q \cdot \text{div}(l) = - \sum_{i \in I_q} V_{i,q}(C_i \cdot \text{div}(l)) = - \sum_{i \in I_q} V_{i,q} \deg_{\mathbb{F}_q}(\mathcal{L}|_{C_{i,q}}).$$

7 Description of the map from the curve to the torsor

The situation is as in Section 2, see (2.12). We describe the morphism $\tilde{j}_b: U \rightarrow T$ in terms of invertible \mathcal{O} -modules on $C \times C^{\text{sm}}$. Since T is the product, over J , of the \mathbb{G}_m -torsors $T_i := (\text{id}, m \cdot \circ \text{tr}_{c_i} \circ f_i)^* P^\times$ this amounts to describing, for each i , the morphism $(\tilde{j}_b)_i: U \rightarrow T_i$. Note that $\text{tr}_{c_i} \circ f_i: J_{\mathbb{Q}} \rightarrow J_{\mathbb{Q}}$ is a morphism of groupschemes composed with a translation, and that all morphisms of schemes $\alpha: J_{\mathbb{Q}} \rightarrow J_{\mathbb{Q}}$ are of this form. From now on we fix one such i and omit it from our notation.

Let $\alpha: J_{\mathbb{Q}} \rightarrow J_{\mathbb{Q}}$ be a morphism of schemes, let \mathcal{L}_α be the pullback of \mathcal{M} (see (6.3.3)) to $C_{\mathbb{Q}} \times C_{\mathbb{Q}}$ via $j_b \times (\alpha \circ j_b)$, and let $T_\alpha := (\text{id}, \alpha)^* \mathcal{M}^\times$ on $J_{\mathbb{Q}}$:

$$(7.1) \quad \begin{array}{ccccc} & & T_\alpha & \xrightarrow{\quad} & \mathcal{M}^\times \\ & & \downarrow & & \swarrow \\ C_{\mathbb{Q}} & \xrightarrow{j_b} & J_{\mathbb{Q}} & \xrightarrow{(\text{id}, \alpha)} & (J \times J)_{\mathbb{Q}} \\ \downarrow \text{diag} & & & & \uparrow j_b \times \text{id} \\ (C \times C)_{\mathbb{Q}} & \xrightarrow{\text{id} \times j_b} & (C \times J)_{\mathbb{Q}} & \xrightarrow{\text{id} \times \alpha} & (C \times J)_{\mathbb{Q}} \\ \uparrow & & & & \swarrow \\ \mathcal{L}_\alpha^\times & \xrightarrow{\quad} & & & \mathcal{L}^{\text{univ}, \times} \end{array}$$

Then $(b, \text{id})^* \mathcal{L}_\alpha = \mathcal{O}_{C_{\mathbb{Q}}}$, \mathcal{L}_α is of degree zero on the fibres of $\text{pr}_2: (C \times C)_{\mathbb{Q}} \rightarrow C_{\mathbb{Q}}$, and: $j_b^* T_\alpha$ is trivial if and only if $\text{diag}^* \mathcal{L}_\alpha$ is trivial. Note that diagram (7.1) without the \mathbb{G}_m -torsors is commutative.

Conversely, let \mathcal{L} be an invertible \mathcal{O} -module on $(C \times C)_{\mathbb{Q}}$, rigidified on $\{b\} \times C_{\mathbb{Q}}$, and of degree 0 on the fibres of $\text{pr}_2: (C \times C)_{\mathbb{Q}} \rightarrow C_{\mathbb{Q}}$. The universal property of $\mathcal{L}^{\text{univ}}$ gives a unique $\beta_{\mathcal{L}}: C_{\mathbb{Q}} \rightarrow J_{\mathbb{Q}}$ such that $(\text{id} \times \beta_{\mathcal{L}})^* \mathcal{L}^{\text{univ}} = \mathcal{L}$ (compatible with rigidification at b). The Albanese property of $j_b: C_{\mathbb{Q}} \rightarrow J_{\mathbb{Q}}$ then gives that $\beta_{\mathcal{L}}$ extends to a unique $\alpha_{\mathcal{L}}: J_{\mathbb{Q}} \rightarrow J_{\mathbb{Q}}$ such that $\alpha_{\mathcal{L}} \circ j_b = \beta_{\mathcal{L}}$. Then $j_b^* T_{\alpha_{\mathcal{L}}}$ is trivial if and only if $\text{diag}^* \mathcal{L}$ is trivial. Now assume that $\text{diag}^* \mathcal{L} = \mathcal{O}_{C_{\mathbb{Q}}}$ (compatible with rigidifications at b), and let

$$(7.2) \quad \ell \in (\text{diag}^* \mathcal{L}^\times)(C_{\mathbb{Q}})$$

correspond to 1. Then $m \cdot \circ \alpha_{\mathcal{L}}$ extends over \mathbb{Z} to $m \cdot \circ \alpha_{\mathcal{L}}: J \rightarrow J^0$, and the restriction of $j_b^*(m \cdot \circ \alpha_{\mathcal{L}})^* \mathcal{M}$ on C^{sm} to U is trivial, giving a lift \tilde{j}_b , unique up to sign:

$$(7.3) \quad \begin{array}{ccccc} & & T_{m \cdot \circ \alpha_{\mathcal{L}}} & \xrightarrow{\quad} & \mathcal{M}^\times \\ & \nearrow \tilde{j}_b & \downarrow & & \downarrow \\ U & \xrightarrow{\quad} & C^{\text{sm}} & \xrightarrow{j_b} & J & \xrightarrow{(\text{id}, m \cdot \circ \alpha_{\mathcal{L}})} & J \times J^0 \end{array}$$

The invertible \mathcal{O} -module \mathcal{L} on $(C \times C)_{\mathbb{Q}}$ with its rigidification of $(b, \text{id})^* \mathcal{L}$, extends uniquely to an invertible \mathcal{O} -module on $(C \times C)_{\mathbb{Z}[1/n]}$, still denoted \mathcal{L} . For S a $\mathbb{Z}[1/n]$ -scheme, d and e in $\mathbb{Z}_{\geq 0}$, $D \in C^{(d)}(S)$ and $E \in C^{(e)}(S)$, we claim that

$$(7.4) \quad \mathcal{M}(\Sigma(D), \alpha_{\mathcal{L}}(\Sigma(E))) = (\text{Norm}_{D/S}(\text{id}, b)^* \mathcal{L})^{\otimes (1-e)} \otimes \text{Norm}_{(D \times E)/S}(\mathcal{L}).$$

To prove this, we may and do assume (finite locally free base change on S) that we have x_i and y_j in $C(S)$, such that $D = \sum_i x_i$ and $E = \sum_j y_j$. Recall that, for $c \in C(S)$, $\beta_{\mathcal{L}}(c)$ in $J(S)$ is $(\text{id}, c)^*\mathcal{L}$ on C_S , with its rigidification at b . Then we have:

$$\begin{aligned}
(7.4.1) \quad \mathcal{M}(\Sigma(D), \alpha_{\mathcal{L}}(\Sigma(E))) &= \mathcal{M}(\alpha_{\mathcal{L}}(\Sigma(E)), \Sigma(D)) \\
&= \mathcal{M}\left(\beta_{\mathcal{L}}(b) + \sum_j (\beta_{\mathcal{L}}(y_j) - \beta_{\mathcal{L}}(b)), \sum_i j_b(x_i)\right) \\
&= \left(\bigotimes_i \mathcal{L}(x_i, b)^{\otimes(1-e)}\right) \otimes \bigotimes_{i,j} \mathcal{L}(x_i, y_j).
\end{aligned}$$

from which the desired equality follows.

For S a $\mathbb{Z}[1/n]$ -scheme and $x \in C(S)$, applying (7.4) with $D = E = x$ gives:

$$(7.5) \quad T_{m \circ \alpha_{\mathcal{L}}}(j_b(x)) = \mathcal{M}^\times(j_b(x), m \cdot \alpha_{\mathcal{L}}(j_b(x))) = \mathcal{L}^{\otimes m}(x, x)^\times = (\mathbb{G}_m)_S,$$

with the last equality coming from the rigidification at b .

Now let \mathcal{L} be any extension of \mathcal{L} with its rigidification of $(b, \text{id})^*\mathcal{L}$ from $(C \times C)_{\mathbb{Z}[1/n]}$ to $C \times U$. For q dividing n , let W_q be the valuation along $U_{\mathbb{F}_q}$ of the rational section ℓ of $\text{diag}^*\mathcal{L}$ on U . Then ℓ , multiplied by the product, over the primes q dividing n , of q^{-W_q} , generates $\text{diag}^*\mathcal{L}$ on U :

$$(7.6) \quad \left(\prod_{q|n} q^{-W_q}\right) \cdot \ell \in (\text{diag}^*\mathcal{L}^\times)(U).$$

There is a unique divisor V on $C \times U$ with support disjoint from $(b, \text{id})U$ and contained in the $(C \times U)_{\mathbb{F}_q}$ with q dividing n , such that

$$(7.7) \quad \mathcal{L}^m := \mathcal{L}^{\otimes m}(V) \quad \text{on } C \times U$$

has multidegree 0 on the fibres of $\text{pr}_2: C \times U \rightarrow U$. Then \mathcal{L}^m is the pullback of $\mathcal{L}^{\text{univ}}$ via $\text{id} \times (m \cdot \circ \alpha_{\mathcal{L}} \circ j_b): C \times U \rightarrow C \times J^0$. Its restriction $\mathcal{L}^m|_{C^{\text{sm}} \times U}$ is then the pullback of \mathcal{M} via $j_b \times (m \cdot \circ \alpha_{\mathcal{L}} \circ j_b): C^{\text{sm}} \times U \rightarrow J \times J^0$, because on $C^{\text{sm}} \times J^0$ the restriction of $\mathcal{L}^{\text{univ}}$ and $(j_b \times \text{id})^*\mathcal{M}$ are equal (both are rigidified after $(b, \text{id})^*$ and equal over $\mathbb{Z}[1/n]$; here we use that, for all $q|n$, $J_{\mathbb{F}_q}^0$ is geometrically connected). Hence, on U we have $j_b^*T_{m \circ \alpha_{\mathcal{L}}} = \text{diag}^*(\mathcal{L}^{\otimes m}(V)^\times)$, compatible with rigidifications at $b \in U(\mathbb{Z}[1/n])$. Our trivialisation \tilde{j}_b on U of $T_{m \circ \alpha_{\mathcal{L}}}$ is therefore a generating section of $\mathcal{L}^{\otimes m}$, multiplied by the product over the q dividing n , of the factors q^{-V_q} , where V_q is the multiplicity in V of the prime divisor $(U \times U)_{\mathbb{F}_q}$. Then, for x and S as in (7.5), we have the following description of \tilde{j}_b :

$$(7.8) \quad \tilde{j}_b(x) = \left(\prod_{q|n} q^{-mW_q - V_q}\right) \cdot \ell^{\otimes m} \quad \text{in } (T_{m \circ \alpha_{\mathcal{L}}}(j_b(x)))(S) = \mathcal{L}^{\otimes m}(x, x)^\times(S).$$

8 An example with genus 2, rank 2, and 14 points

The example that we are going to treat is the quotient of the modular curve $X_0(129)$ by the action of the group of order 4 generated by the Atkin-Lehner involutions w_3 and w_{43} . An

equation for this quotient is given in the table in [14], and Magma has shown that that equation and the equations below give isomorphic curves over \mathbb{Q} .

Let C_0 be the curve over \mathbb{Z} obtained from the following closed subschemes of $\mathbb{A}_{\mathbb{Z}}^2$

$$\begin{aligned} V_1 : \quad y^2 + y &= x^6 - 3x^5 + x^4 + 3x^3 - x^2 - x, \\ V_2 : \quad w^2 + z^3w &= 1 - 3z + z^2 + 3z^3 - z^4 - z^5 \end{aligned}$$

by glueing the open subset of V_1 where x is invertible with the open subset of V_2 where z is invertible using the identifications $z = 1/x$, $w = y/x^3$. The scheme C_0 can be also described as a subscheme of the line bundle \mathcal{L}_3 associated to the invertible \mathcal{O} -module $\mathcal{O}_{\mathbb{P}_{\mathbb{Z}}^1}(3)$ on $\mathbb{P}_{\mathbb{Z}}^1$ with homogeneous coordinates X, Z : the map $\mathcal{O}_{\mathbb{P}_{\mathbb{Z}}^1}(3) \rightarrow \mathcal{O}_{\mathbb{P}_{\mathbb{Z}}^1}(6)$ sending a section Y to $Y \otimes Y + Z^3 \otimes Y$ induces a map φ from \mathcal{L}_3 to the line bundle \mathcal{L}_6 associated to $\mathcal{O}(6)$; then C_0 is isomorphic to the inverse image by φ of the section $s := X^6 - 3X^5Z + X^4Z^2 + 3X^3Z^3 - X^2Z^4 - XZ^5$ of \mathcal{L}_6 and since the map φ is finite of degree 2 then C_0 is finite of degree 2 over $\mathbb{P}_{\mathbb{Z}}^1$. Hence C_0 is proper over \mathbb{Z} and it is moreover smooth over $\mathbb{Z}[1/n]$ with $n = 3 \cdot 43$. The generic fiber of C_0 is a curve of genus $g = 2$, labeled 5547.b.16641.1 on www.lmfdb.org. The only point where C_0 is not regular is the point $P_0 = (3, x - 2, y - 1)$ contained in V_1 and the blow up C of C_0 in P_0 is regular.

In the rest of the article we apply our geometric method to the curve C and we prove that $C(\mathbb{Z})$ contains exactly 14 elements. We use the same notation as in Sections 2 and 4.

The fiber $C_{\mathbb{F}_{43}}$ is absolutely irreducible while $C_{\mathbb{F}_3}$ is the union of two geometrically irreducible curves, a curve of genus 0 that lies above the point P_0 and that we call K_0 , and a curve of genus 1 that we call K_1 . We define $U_0 := C \setminus K_1$ and $U_1 := C \setminus K_0$ so that $C(\mathbb{Z}) = C^{\text{sm}}(\mathbb{Z}) = U_0(\mathbb{Z}) \cup U_1(\mathbb{Z})$ and both U_0 and U_1 satisfy the hypothesis of U in Section 2. We have $K_0 \cdot K_1 = 2$ and consequently the self-intersections of K_0 and K_1 are both equal to -2 . We deduce that all the fibers of J over \mathbb{Z} are connected except for $J_{\mathbb{F}_3}$ which has group of connected components equal to $\mathbb{Z}/2\mathbb{Z}$. Hence $m = 2$.

The automorphism group of C is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^2$, generated by the automorphisms ι and η lifting the extension to C_0 of

$$\iota, \eta: V_1 \longrightarrow V_1, \quad \iota: (x, y) \longmapsto (x, -1 - y), \quad \eta: (x, y) \longmapsto (1 - x, -1 - y).$$

The quotients $E_1 := C_{\mathbb{Q}}/\eta$ and $E_2 := C_{\mathbb{Q}}/(\iota \circ \eta)$ are curves of genus 1 and the two projections $C \rightarrow E_i$ induce an isogeny $J \rightarrow \text{Pic}^0(E_1) \times \text{Pic}^0(E_2)$. The elliptic curves $\text{Pic}^0(E_i)$ are not isogenous and $\rho = 2$.

8.1 The torsor on the jacobian

Let $\infty, \infty_- \in C(\mathbb{Z})$ be the lifts of $(0, 1), (0, -1) \in V_2(\mathbb{Z}) \subset C_0(\mathbb{Z})$ and let us fix the base point $b = \infty$ in $C(\mathbb{Z})$. Following Section 7 we describe a \mathbb{G}_m -torsor $T \rightarrow J$ and maps $\widetilde{j}_{b,i}: U_i \rightarrow T$ using invertible \mathcal{O} -modules on $C \times C^{\text{sm}}$. Let Γ_{η} denote the graph of the map $\eta: C \rightarrow C$. Over $C_{\mathbb{Q}} \times C_{\mathbb{Q}}$ we consider

$$\mathcal{L} := \mathcal{O}_{C_{\mathbb{Q}} \times C_{\mathbb{Q}}}(\Gamma_{\eta, \mathbb{Q}} - \infty_- \times C_{\mathbb{Q}} - C_{\mathbb{Q}} \times \infty)$$

trivialised on $b \times C_{\mathbb{Q}}$ through the section

$$l_b := 2 \quad \text{in } ((b, \text{id})^* \mathcal{L})(C_{\mathbb{Q}}) = \mathcal{O}_{C_{\mathbb{Q}}}(\eta(b) - b)(C_{\mathbb{Q}}) = \mathcal{O}_{C_{\mathbb{Q}}}(C_{\mathbb{Q}}).$$

Notice that $(\text{id}, b)^* \mathcal{L}$ has degree 0 hence there exists a morphism of schemes $\alpha_{\mathcal{L}}: J_{\mathbb{Q}} \rightarrow J_{\mathbb{Q}}$ such that \mathcal{L} , with its rigidification at $b \times C_{\mathbb{Q}}$, is the pullback of \mathcal{M} via $j_b \times (\alpha_{\mathcal{L}} \circ j_b)$. For every $\overline{\mathbb{Q}}$ -point Q on $C_{\mathbb{Q}}$ the invertible $\mathcal{O}_{C_{\overline{\mathbb{Q}}}}$ -module $(\text{id}, Q)^* \mathcal{L}$ is isomorphic to $\mathcal{O}_{C_{\overline{\mathbb{Q}}}}(\eta(Q) - \infty_-)$ hence

$$\alpha = \text{tr}_c \circ f, \quad \text{with } f = \eta_* \text{ and } c = [D_0], D_0 := \infty - \infty_-.$$

When restricted to the diagonal \mathcal{L} is trivial since, compatibly with the trivialisation at (b, b) ,

$$\text{diag}^* \mathcal{L} = \mathcal{O}_{C_{\mathbb{Q}}}(\infty_- + \infty - \infty_- - \infty) = \mathcal{O}_{C_{\mathbb{Q}}} = l \cdot \mathcal{O}_{C_{\mathbb{Q}}} \quad \text{with } l := 1.$$

Following Section 7 we choose the extension of \mathcal{L} over $C \times C^{\text{sm}}$

$$\mathcal{L} := \mathcal{O}(\Gamma_{\eta}|_{C \times C^{\text{sm}}} - \infty_- \times C^{\text{sm}} - C \times \infty)$$

trivialised along $b \times C^{\text{sm}}$ through the section $l_b = 2$ (the points ∞_- and b have a simple intersection over the prime 2). Following (7.5) the torsor $T := T_{m \circ \alpha_{\mathcal{L}}}$ on J satisfies, for S a $\mathbb{Z}[1/n]$ -scheme and x in $C(S)$, using the trivialisation given by l and l_b

$$\begin{aligned} (8.1.1) \quad T(j_b(x)) &= \mathcal{M}^{\times}(j_b(x), m \cdot \alpha_{\mathcal{L}}(j_b(x))) = \mathcal{M}^{\times}(j_b(x), (\text{id}, x)^* \mathcal{L}^{\otimes m}) \\ &= x^*(\text{id}, x)^* \mathcal{L}^{\otimes m, \times} \otimes b^*(\text{id}, x)^* \mathcal{L}^{\otimes -m, \times} \\ &= \mathcal{L}^{\otimes m, \times}(x, x) \otimes \mathcal{L}^{\otimes m, \times}(b, x)^{-1} = \mathcal{L}^{\otimes m, \times}(x, x) = \mathcal{O}_S^{\times}. \end{aligned}$$

Since l generates $\text{diag}^*(\mathcal{L})$ on the whole C^{sm} then, we have $W_3 = W_{43} = 0$ in (7.6) when computing both $\widetilde{j}_{b,0}$ and $\widetilde{j}_{b,1}$.

The invertible \mathcal{O} -module $\mathcal{L}^{\otimes m}$ has multidegree 0 over all the fibers $C \times U_1 \rightarrow U_1$, hence in order to compute $\widetilde{j}_{b,1}$ we must take $V = 0$ in (7.7), giving $V_3 = V_{43} = 0$. Hence for S and x as in (8.1.1), assuming moreover that 2 is invertible on S ,

$$(8.1.2) \quad \widetilde{j}_{b,1}(x) = l^2 \otimes l_b^{-2} = \frac{1}{4}(x^*1) \otimes (b^*1)^{-1} \quad \text{in}$$

$T(j_b(x)) = x^*(\text{id}, x)^* \mathcal{L}^{\otimes m, \times} \otimes b^*(\text{id}, x)^* \mathcal{L}^{\otimes -m, \times} = x^* \mathcal{O}_{C_S}(\eta(x) - \infty_-)^{\times} \otimes b^* \mathcal{O}_{C_S}(\eta(x) - \infty_-)^{\times}$, where the last equality in (8.1.2) makes sense if the image of x is disjoint from ∞, ∞_- in C_S .

The restriction $\mathcal{L}^{\otimes m}$ to $C \times U_0$ has multidegree 0 over all the fibers $C \times U_0 \rightarrow U_0$ of characteristic not 3, while if we consider a fiber of characteristic 3 it has degree 2 over K_0 and degree -2 over K_1 . Hence for computing $\widetilde{j}_{b,0}$ we take $V = K_0 \times (K_0 \cap U_0)$ in (7.7) giving $V_{43} = 0, V_3 = 1$. Hence for S and x as in (8.1.1), assuming moreover that 2 is invertible on S ,

$$(8.1.3) \quad \widetilde{j}_{b,0}(x) = \frac{1}{3}l^2 \otimes l_b^{-2} = \frac{1}{12}(x^*1) \otimes (b^*1)^{-1} \quad \text{in}$$

$T(j_b(x)) = x^*(\text{id}, x)^* \mathcal{L}^{\otimes m, \times} \otimes b^*(\text{id}, x)^* \mathcal{L}^{\otimes -m, \times} = x^* \mathcal{O}_{C_S}(\eta(x) - \infty_-)^{\times} \otimes b^* \mathcal{O}_{C_S}(\eta(x) - \infty_-)^{\times}$, where the last equality in (8.1.3) makes sense if the image of x is disjoint from ∞, ∞_- in C_S .

8.2 Some integral points on the biextension

On C_0 we have the following integral points that lift uniquely to elements of $C(\mathbb{Z})$

$$\begin{aligned} \infty &= (0, 1), & \infty_- &:= (0, -1) \quad \text{in } V_2(\mathbb{Z}), \\ \alpha &:= (1, 0), & \beta &:= \eta(\alpha) = (0, -1), & \gamma &:= (2, 1), & \delta &:= \eta(\gamma) = (-1, -2) \quad \text{in } V_1(\mathbb{Z}). \end{aligned}$$

Computations in Magma confirm that $J(\mathbb{Z})$ is a free \mathbb{Z} -module of rank $r = 2$ generated by

$$G_1 := \gamma - \alpha, \quad G_2 := \alpha + \infty_- - 2\infty.$$

The points in $T(\mathbb{Z})$ are a subset of points of $\mathcal{M}^\times(\mathbb{Z})$ that can be constructed, using the two group laws, from the points in $\mathcal{M}^\times(G_i, m \cdot f(G_j))(\mathbb{Z})$ and $\mathcal{M}^\times(G_i, m \cdot D_0)(\mathbb{Z})$ for $i, j \in \{1, 2\}$. Let us compute in detail $\mathcal{M}^\times(G_1, m \cdot f(G_1))(\mathbb{Z})$. As explained in Section 6.9

$$\begin{aligned} \mathcal{M}(G_1, m \cdot f(G_1))^\times &= \mathcal{M}^\times(\gamma - \alpha, 2\delta - 2\beta) \\ &= 3^{e_3} 43^{e_{43}} \cdot \text{Norm}_{(2\delta)/\mathbb{Z}}(\mathcal{O}_C(\gamma - \alpha)) \otimes \text{Norm}_{(2\beta)/\mathbb{Z}}(\mathcal{O}_C(\gamma - \alpha))^{-1} \\ &= 3^{e_3} 43^{e_{43}} \cdot (2\delta - 2\beta)^* \mathcal{O}_C(\gamma - \alpha) \end{aligned}$$

where, given a scheme S , an invertible \mathcal{O} -module \mathcal{L} on C_S and a divisor $D_+ - D_- = \sum_i n_i P_i$ on C_S that is sum of S -points, we define the invertible \mathcal{O}_S -module

$$\left(\sum_i n_i P_i \right)^* \mathcal{L} := \bigotimes_i P_i^* \mathcal{L}^{n_i} = \text{Norm}_{D_+/S}(\mathcal{L}) \otimes \text{Norm}_{D_-/S}(\mathcal{L})^{-1}.$$

Since $C_{\mathbb{F}_{43}}$ is irreducible then $2f(G_1)$ has already multidegree 0 over 43, hence $e_{43} = 0$. If we look at $C_{\mathbb{F}_3}$ then $2f(G_1)$ does not have multidegree 0, while $2f(G_1) + K_0$ has multidegree 0; hence by (6.9.12)

$$e_3 = -\deg_{\mathbb{F}_3} \mathcal{O}_C(\gamma - \alpha)|_{K_0} = -1.$$

Notice that over $\mathbb{Z}[\frac{1}{2}]$ the divisor G_1 is disjoint from β and δ (to see that it is disjoint from $\delta = (-1, -2, 1)$ over the prime 3 one needs to look at local equations of the blow up) thus $\beta^* \mathcal{O}_C(\gamma - \alpha)$ and $\delta^* \mathcal{O}_C(\gamma - \alpha)$ are generated by $\beta^* 1$ and $\delta^* 1$ over $\mathbb{Z}[\frac{1}{2}]$. Thus there are integers e_β, e_δ such that $\beta^* \mathcal{O}_C(\gamma - \alpha)$ and $\delta^* \mathcal{O}_C(\gamma - \alpha)$ are generated by $\beta^* 2^{e_\beta}$ and $\delta^* 2^{e_\delta}$ over \mathbb{Z} . Looking at the intersections between β, γ, α and δ we compute that $e_\beta = -1$ $e_\delta = 1$ hence

$$\begin{aligned} \mathcal{M}(G_1, m \cdot f(G_1)) &= 3^{-1} \cdot (\delta^* 2)^2 \otimes (\beta^* 2^{-1})^{-2} \cdot \mathbb{Z} = 2^4 \cdot 3^{-1} \cdot (\delta^* 1)^2 \otimes (\beta^* 1) \cdot \mathbb{Z} \quad \text{and} \\ Q_{1,1} &:= \pm 2^4 \cdot 3^{-1} \cdot (\delta^* 1)^2 \otimes (\beta^* 1)^{-2} \in \mathcal{M}_{G_1, m \cdot f(G_1)}^\times(\mathbb{Z}). \end{aligned}$$

With analogous computations we see that

$$\begin{array}{ll} Q_{2,1} := 2^{-2} \cdot (\delta^* 1)^2 \otimes (\beta^* 1)^{-2} & \text{generates } \mathcal{M}_{G_2, m \cdot f(G_1)} \\ Q_{1,2} := 2^{-2} \cdot (\beta^* 1)^2 \otimes (\infty^* 1)^2 \otimes (\infty^* 1)^{-4} & \text{generates } \mathcal{M}_{G_1, m \cdot f(G_2)} \\ Q_{2,2} := 2^{18} \cdot (\beta^* 1)^2 \otimes (\infty^* x)^2 \otimes (\infty^* z^2)^{-4} & \text{generates } \mathcal{M}_{G_2, m \cdot f(G_2)} \\ Q_{1,2} := (\infty^* 1)^2 \otimes (\infty^* 1)^{-2} & \text{generates } \mathcal{M}_{G_1, m \cdot D_0} \\ Q_{2,0} := 2^{-12} \cdot (\infty^* z^2)^2 \otimes (\infty^* x)^{-2} & \text{generates } \mathcal{M}_{G_2, m \cdot D_0}. \end{array}$$

8.3 Some residue disks of the biextension

Let p be a prime of good reduction for C . Given the divisors

$$D := \alpha - \infty, \quad E := 2\beta - 2\infty_- = (m \circ \text{tr}_c \circ \eta_*)(D) \quad \text{in } \text{Div}(C_{\mathbb{Z}/p^2})$$

we apply Section 6.6 and we give parameters on the residue disks in $\mathcal{M}^\times(\mathbb{Z}/p^2)_{\overline{D}, \overline{E}}$ and $T(\mathbb{Z}/p^2)_{\overline{D}}$, with $\overline{D}, \overline{E}$ the images of D, E in $\text{Div}(C_{\mathbb{F}_p})$.

We choose the ‘‘base points’’ $b_1 = \alpha, b_2 = \infty, b_3 = \beta, b_4 = \infty$, so that $b_1 \neq b_2, b_3 \neq b_4$ and $h^0(C_{\mathbb{F}_p}, b_1 + b_2) = h^0(C_{\mathbb{F}_p}, b_3 + b_4) = 1$. As in Section 6.6 we define $x_\alpha = x - 1, x_\infty = z, x_\beta = x$ and $x_{D, \beta} = x_{D, \infty_-} = 1, x_{D, \infty} = z^{-1}$. For Q in $\{\infty, \beta, \alpha\}$ and $a \in \mathbb{F}_p$ let Q_a be the unique \mathbb{Z}/p^2 -point of C that is congruent to Q modulo p and such that $x_Q(Q_a) = ap \in \mathbb{Z}/p^2$. We have the bijections

$$\begin{aligned} \mathbb{F}_p^2 &\longrightarrow J(\mathbb{Z}/p^2)_{\overline{D}}, & \lambda &\longmapsto D_\lambda := D + \alpha_{\lambda_1} - \alpha + \infty_{\lambda_2} - \infty = \alpha_{\lambda_1} + \infty_{\lambda_2} - 2\infty \\ \mathbb{F}_p^2 &\longrightarrow J(\mathbb{Z}/p^2)_{\overline{E}}, & \mu &\longmapsto E_\mu := E + \beta_{\mu_1} - \beta + \infty_{\mu_2} - \infty = \beta + \beta_{\mu_1} + \infty_{\mu_2} - \infty - 2\infty_-. \end{aligned}$$

Following (6.6.6) for $\lambda, \mu \in \mathbb{F}_p^2$ we define

$$s_{D,E}(\lambda, \mu) := (\beta^* 1) \otimes (\beta_{\mu_1}^* 1) \otimes \left(\infty_{\mu_2}^* \frac{z^2}{z - \lambda_2 p} \right) \otimes \left(\infty^* \frac{z^2}{z - \lambda_2 p} \right)^{-1} \otimes (\infty_-^* 1)^{-2}$$

that, by Proposition 6.3.2 and Remark 6.3.12, generates $E_\mu^* \mathcal{O}_{C_{\mathbb{Z}/p^2}}(D_\lambda) = \mathcal{M}_{D_\lambda, E_\mu}$. The points in $\mathcal{M}^\times(\mathbb{F}_p)$ projecting to $(\overline{D}, \overline{E})$ are in bijection with the elements ξ in \mathbb{F}_p^\times and are exactly the points $\xi \cdot s_{D,E}(0, 0)$. Using $(\mathbb{Z}/p^2)^\times = \mathbb{F}_p^\times \times (1 + p\mathbb{F}_p)$, for each $\xi \in \mathbb{F}_p^\times$ we parametrise the residue disk of $\xi \cdot s_{D,E}(0, 0)$ using bijection (6.6.7)

$$\mathbb{F}_p^5 \longrightarrow \mathcal{M}^\times(\mathbb{Z}/p^2)_{\xi \cdot s_{D,E}(0,0)}, \quad (\lambda_1, \lambda_2, \mu_1, \mu_2, \tau) \longmapsto (1 + p\tau)\xi \cdot s_{D,E}((\lambda_1, \lambda_2), (\mu_1, \mu_2)).$$

Since $(m \circ \text{tr}_c \circ f)(D_\lambda) = E_{-2\lambda}$ then we have

$$T(\mathbb{Z}/p^2)_{\overline{D}} = \bigcup_{\lambda \in \mathbb{F}_p^2} T_{D_\lambda}(\mathbb{Z}/p^2) = \bigcup_{\lambda \in \mathbb{F}_p^2} \mathcal{M}_{D_\lambda, E_{-2\lambda}}^\times(\mathbb{Z}/p^2).$$

As ξ varies in \mathbb{F}_p^\times the point $\xi \cdot s_{D,E}(0, 0)$ varies in all the points in $\mathcal{M}^\times(\mathbb{F}_p)$ projecting to $(\overline{D}, \overline{E})$ and we have the following bijection induced by parameters in $\xi \cdot s_{D,E}(0, 0)$

$$(8.3.1) \quad \mathbb{F}_p^3 \longrightarrow T(\mathbb{Z}/p^2)_{\xi \cdot s_{D,E}(0,0)}, \quad (\lambda_1, \lambda_2, \tau) \longmapsto (1 + \tau p) \cdot \xi \cdot s_{D,E}((\lambda_1, \lambda_2), (-2\lambda_1, -2\lambda_2)).$$

If we apply (8.1.2) and (8.1.3) to $Q = \alpha_\lambda$ and we use the symmetry of the Poincaré torsor explained in Proposition 6.3.2 and made explicit in Section 6.5 we obtain the following description of $\widetilde{j}_{b,i}$ on $C(\mathbb{Z}/p^2)_{\alpha_{\mathbb{F}_p}}$ when $p \neq 2$

$$\widetilde{j}_{b,1}(\alpha_\lambda) = (1/4) \cdot s_{D,E}((\lambda, 0), (-2\lambda, 0)), \quad \widetilde{j}_{b,0}(Q) = (1/12) \cdot s_{D,E}((\lambda, 0), (-2\lambda, 0)).$$

If $p = 5$ then 18 and -1 are $(p - 1)$ -th roots of unity in $(\mathbb{Z}/p^2)^\times$, thus $1/4 = (-1)(1 + p)$ and $1/12 = 3(1 + 2p)$ in $(\mathbb{Z}/p^2)^\times = \mathbb{F}_p^\times \times (1 + p\mathbb{F}_p)$, hence

$$(8.3.2) \quad \widetilde{j}_{b,1}(\alpha_\lambda) = -(1 + p) \cdot s_{D,E}((\lambda, 0), (-2\lambda, 0)), \quad \widetilde{j}_{b,0}(Q) = 3 \cdot (1 + 2p) \cdot s_{D,E}((\lambda, 0), (-2\lambda, 0)).$$

Since it is useful for computing the map $\kappa_{\mathbb{Z}}$ in the residue disks of $T(\mathbb{Z}/p^2)$ projecting to \overline{D} , let us apply the recipe in Section 6.6 also to the residue disks of $\mathcal{M}^\times(\mathbb{Z}/p^2)$ lying over $(\overline{D}, 0)$, $(0, \overline{E})$ and $(0, 0)$. Hence for $\lambda, \mu \in \mathbb{F}_p^2$ we define the divisors on $C_{\mathbb{Z}/p^2}$

$$D_\lambda^0 := \alpha_{\lambda_1} - \alpha + \infty_{\lambda_2} - \infty, \quad E_\mu^0 := \beta_{\mu_1} - \beta + \infty_{\mu_2} - \infty$$

and the sections

$$\begin{aligned} s_{D,0}(\lambda, \mu) &:= (\beta_{\mu_1}^* 1) \otimes (\infty_{\mu_2}^* \frac{z^2}{z-\lambda_2 p}) \otimes (\beta^* 1)^{-1} \otimes (\infty^* \frac{z^2}{z-\lambda_2 p})^{-1} && \text{in } \mathcal{M}^\times(D_\lambda, E_\mu^0)(\mathbb{Z}/p^2) \\ s_{0,E}(\lambda, \mu) &:= (\beta^* 1) \otimes (\beta_{\mu_1}^* 1) \otimes (\infty_{\mu_2}^* \frac{z}{z-\lambda_2 p}) \otimes (\infty^* \frac{z}{z-\lambda_2 p})^{-1} \otimes (\infty^* 1)^{-2} && \text{in } \mathcal{M}^\times(D_\lambda^0, E_\mu)(\mathbb{Z}/p^2) \\ s_{0,0}(\lambda, \mu) &:= (\beta_{\mu_1}^* 1) \otimes (\infty_{\mu_2}^* \frac{z}{z-\lambda_2 p}) \otimes (\beta^* 1)^{-1} \otimes (\infty^* \frac{z}{z-\lambda_2 p})^{-1} && \text{in } \mathcal{M}^\times(D_\lambda^0, E_\mu^0)(\mathbb{Z}/p^2). \end{aligned}$$

8.4 Geometry mod p of integral points

From now on $p = 5$. Let $\overline{\alpha} \in C(\mathbb{Z}/p^2)$ be the image of $\alpha \in C(\mathbb{Z})$. In this subsection we compute the composition $\overline{\kappa}: \mathbb{Z}^2 \rightarrow T(\mathbb{Z}/p^2)_{\widetilde{j_{b,1}(\overline{\alpha})}}$ of the map $\kappa_{\mathbb{Z}}: \mathbb{Z}^2 \rightarrow T(\mathbb{Z}_p)_{\widetilde{j_{b,1}(\overline{\alpha})}}$ in (4.9) and the reduction map $T(\mathbb{Z}_p)_{\widetilde{j_{b,1}(\overline{\alpha})}} \rightarrow T(\mathbb{Z}/p^2)_{\widetilde{j_{b,1}(\overline{\alpha})}}$. With a suitable choice of parameters in $\mathcal{O}_{T, \widetilde{j_{b,1}(\overline{\alpha})}}$, the map $\kappa_{\mathbb{Z}}$ is described by integral convergent power series $\kappa_1, \kappa_2, \kappa_3 \in \mathbb{Z}_p\langle z_1, z_2 \rangle$ and $\overline{\kappa}$, composed with the inverse of the parametrization (8.3.1), is given the images $\overline{\kappa}_1, \overline{\kappa}_2, \overline{\kappa}_3$ of $\kappa_1, \kappa_2, \kappa_3$ in $\mathbb{F}_p[z_1, z_2]$.

The divisor $j_b(\overline{\alpha})$ is equal to the image of

$$\widetilde{G}_t := e_{0,1}G_1 + e_{0,2}G_2 \text{ with } e_{0,1} := 6, e_{0,2} := 3$$

in $J(\mathbb{F}_p)$ and

$$\tilde{t} := Q_{1,0}^6 \otimes Q_{2,0}^3 \otimes Q_{1,1}^{6 \cdot 6} \otimes Q_{1,2}^{6 \cdot 3} \otimes Q_{2,1}^{3 \cdot 6} \otimes Q_{2,2}^{3 \cdot 3} \text{ in } \mathcal{M}^\times(\widetilde{D}_1, m \cdot (D_0 + \eta_* \widetilde{G}_t))(\mathbb{Z})$$

is a lift of $\widetilde{j_{b,1}(\overline{\alpha})}$. The kernel of $J(\mathbb{Z}) \rightarrow J(\mathbb{F}_p)$ is a free \mathbb{Z} -module generated by

$$\widetilde{G}_1 := e_{1,1}G_1 + e_{1,2}G_2, \quad \widetilde{G}_2 := e_{2,1}G_1 + e_{2,2}G_2, \text{ with } e_{1,1} := 16, e_{1,2} := 2, e_{2,1} := 0, e_{2,2} := 5.$$

Let $\widetilde{G}_{t,2}$ be the divisor $m(D_0 + \eta_*(\widetilde{G}_t))$ representing $(m \cdot \text{otr}_{c \circ f})(\widetilde{G}_t) \in J^0(\mathbb{Z})$. Following (4.1) for $i, j \in \{1, 2\}$ we define

$$\begin{array}{ccc} P_{i,j} := \bigotimes_{m,l=1}^2 Q_{l,m}^{e_{i,l} \cdot e_{j,m}} & R_{i,\tilde{t}} := \bigotimes_{l=1}^2 Q_{l,0}^{e_{i,l}} \otimes \bigotimes_{m,l=1}^2 Q_{l,m}^{e_{i,l} \cdot e_{0,m}} & S_{i,j} := \bigotimes_{m,l=1}^2 Q_{l,m}^{e_{0,l} \cdot e_{j,m}} \\ \downarrow & \downarrow & \downarrow \\ (\widetilde{G}_i, f(m\widetilde{G}_j)) & (\widetilde{G}_i, \widetilde{G}_{t,2}) & (\widetilde{G}_t, f(m\widetilde{G}_j)). \end{array}$$

Computations in $C_{\mathbb{Z}/p^2}$ show the following linear equivalences of divisors

$$\widetilde{G}_t \sim D_{0,3}, \quad \widetilde{G}_1 \sim D_{4,0}^0, \quad \widetilde{G}_2 \sim D_{0,3}^0$$

and applying Section 6.4 and the functoriality of the norm we compute

(8.4.1)

$$\begin{aligned}
P_{1,1} &= (1+4p) \cdot s_{0,0}((4,0), (2,0)) && \text{in } \mathcal{M}^\times(\widetilde{G}_1, \widetilde{G}_1)(\mathbb{Z}/p^2) = \mathcal{M}^\times(D_{4,0}^0, E_{2,0}^0)(\mathbb{Z}/p^2), \\
P_{1,2} &= (1+4p) \cdot s_{0,0}((4,0), (0,4)) && \text{in } \mathcal{M}^\times(\widetilde{G}_1, \widetilde{G}_2)(\mathbb{Z}/p^2) = \mathcal{M}^\times(D_{4,0}^0, E_{0,4}^0)(\mathbb{Z}/p^2), \\
P_{2,1} &= (1+4p) \cdot s_{0,0}((0,3), (2,0)) && \text{in } \mathcal{M}^\times(\widetilde{G}_2, \widetilde{G}_1)(\mathbb{Z}/p^2) = \mathcal{M}^\times(D_{0,3}^0, E_{2,0}^0)(\mathbb{Z}/p^2), \\
P_{2,2} &= (-1) \cdot (1+2p) \cdot s_{0,0}((0,3), (0,4)) && \text{in } \mathcal{M}^\times(\widetilde{G}_2, \widetilde{G}_2)(\mathbb{Z}/p^2) = \mathcal{M}^\times(D_{0,3}^0, E_{0,4}^0)(\mathbb{Z}/p^2), \\
R_{1,\tilde{t}} &= s_{0,E}((4,0), (0,4)) && \text{in } \mathcal{M}^\times(\widetilde{G}_1, \widetilde{G}_{t,2})(\mathbb{Z}/p^2) = \mathcal{M}^\times(D_{4,0}^0, E_{0,4}^0)(\mathbb{Z}/p^2), \\
R_{2,\tilde{t}} &= (1+4p) \cdot s_{0,E}((0,3), (0,4)) && \text{in } \mathcal{M}^\times(\widetilde{G}_2, \widetilde{G}_{t,2})(\mathbb{Z}/p^2) = \mathcal{M}^\times(D_{0,3}^0, E_{0,4}^0)(\mathbb{Z}/p^2), \\
S_{\tilde{t},1} &= s_{D,0}((0,3), (2,0)) && \text{in } \mathcal{M}^\times(\widetilde{G}_t, \widetilde{G}_1)(\mathbb{Z}/p^2) = \mathcal{M}^\times(D_{0,3}, E_{2,0}^0)(\mathbb{Z}/p^2), \\
S_{\tilde{t},2} &= (-1)(1+4p) \cdot s_{D,0}((0,3), (0,4)) && \text{in } \mathcal{M}^\times(\widetilde{G}_t, \widetilde{G}_2)(\mathbb{Z}/p^2) = \mathcal{M}^\times(D_{0,3}, E_{0,4}^0)(\mathbb{Z}/p^2), \\
\tilde{t} &= (-1) \cdot (1+2p) \cdot s_{D,E}((0,3), (0,4)) && \text{in } \mathcal{M}^\times(\widetilde{G}_t, \widetilde{G}_{t,2})(\mathbb{Z}/p^2) = \mathcal{M}^\times(D_{0,3}, E_{0,4}^0)(\mathbb{Z}/p^2).
\end{aligned}$$

We now show these computations in the cases of \widetilde{G}_t and \tilde{t} . The Riemann-Roch space relative to the divisor $\widetilde{G}_t + \infty + \alpha - D$ on $C_{\mathbb{Z}/p^2}$ is generated by the inverse of the rational function

$$h_1 := \frac{x^9 - 5x^8 - 2x^7 + 7x^6 - 9x^5 - 5x^4 + 14x^3 + 7x^2 + 13x + 1 + (x^6 + 9x^5 - 5x^4 + 15x^3 - 5x^2 + 4x + 14)y}{15x^5 - x^4 + 4x^3 + 19x^2 + 4x + 9}$$

and indeed

$$\text{div}(h_1) = \widetilde{G}_t - D_{0,3} = (6\gamma + 3\infty_- - 3\alpha - 6\infty) - (\alpha + \infty_3 - 2\infty) \quad \text{in } \text{Div}(C_{\mathbb{Z}/p^2}).$$

Hence multiplication by h_1 gives an isomorphism $\mathcal{O}_{C_{\mathbb{Z}/p^2}}(\widetilde{G}_t) \rightarrow \mathcal{O}_{C_{\mathbb{Z}/p^2}}(D_{0,3})$ and by functoriality of the norm we get

$$\begin{aligned}
\delta^* \mathcal{O}_C(\widetilde{G}_t) &\rightarrow \delta^* \mathcal{O}_{C_{\mathbb{Z}/p^2}}(D_{0,3}), && \delta^* 1 \mapsto \delta^*(h_1) = h_1(\delta) \cdot \delta^* 1 = 12 \cdot \delta^* 1, \\
\beta^* \mathcal{O}_C(\widetilde{G}_t) &\rightarrow \beta^* \mathcal{O}_{C_{\mathbb{Z}/p^2}}(D_{0,3}), && \beta^* 1 \mapsto \beta^*(h_1) = h_1(\beta) \cdot \beta^* 1 = 18 \cdot \beta^* 1, \\
\infty^* \mathcal{O}_C(\widetilde{G}_t) &\rightarrow \infty^* \mathcal{O}_{C_{\mathbb{Z}/p^2}}(D_{0,3}), && \infty^* z^6 \mapsto \infty^*(z^6 h_1) = 13 \cdot \infty^* \frac{z^2}{z-3p}, \\
\infty_-^* \mathcal{O}_C(\widetilde{G}_t) &\rightarrow \infty_-^* \mathcal{O}_{C_{\mathbb{Z}/p^2}}(D_{0,3}), && \infty_-^* z^{-3} \mapsto \infty_-^*(z^{-3} h_1) = (z^{-3} h_1)(\infty_-) \cdot \infty_-^* 1 = 6 \cdot \infty_-^* 1.
\end{aligned}$$

Since $\widetilde{G}_{t,2} = 12\delta + 4\infty_- - 6\beta - 10\infty$, the above isomorphisms, tensored with the exponents, give the canonical isomorphism

$$(8.4.2) \quad \mathcal{M}(\widetilde{G}_t, \widetilde{G}_{t,2}) = \widetilde{G}_{t,2}^* \mathcal{O}_{C_{\mathbb{Z}/p^2}}(\widetilde{G}_t) \rightarrow \widetilde{G}_{t,2}^* \mathcal{O}_{C_{\mathbb{Z}/p^2}}(D_{0,3}) = \mathcal{M}(D_{0,3}, \widetilde{G}_{t,2})$$

$$\tilde{t} = 14 \cdot (\delta^* 1)^{12} \otimes (\beta^* 1)^{-6} \otimes (\infty^* z^6)^{-10} \otimes (\infty_-^* z^{-3})^4 \mapsto 14 \cdot (\delta^* 1)^{12} \otimes (\beta^* 1)^{-6} \otimes (\infty^* \frac{z^2}{z-3p})^{-10} \otimes (\infty_-^* 1)^4.$$

The Riemann-Roch space relative to the divisor $\widetilde{G}_{t,2} + \infty + \alpha - E$ on $C_{\mathbb{Z}/p^2}$ is generated by the inverse of the rational function

$$\begin{aligned}
h_2 := & \frac{x^{17} - 8x^{16} + x^{15} - 4x^{14} + 7x^{13} + 4x^{12} + 12x^{11} + x^{10} + 2x^9 - 5x^8 + x^7 + 3x^6 + 12x^5 - 6x^4 - 6x^3 + 4x^2 - 6}{20x^8 - 6x^7} \\
& + \frac{10x^2 + (x^{15} + 6x^{14} - 5x^{13} - x^{12} - 2x^{11} + 14x^{10} - 4x^9 + 14x^8 + 3x^7 + 8x^6 - 6x^5 - 3x^4 + 4x^3 + 13x^2 - x - 7)y}{20x^9 - 6x^8}
\end{aligned}$$

and indeed

$$\operatorname{div}(h_2) = \widetilde{G}_{t,2} - E_{0,4} = (12\delta + 4\infty_- - 6\beta - 10\infty) - (2\beta + \infty_4 - \infty - \infty_-) \quad \text{in } \operatorname{Div}(C_{\mathbb{Z}/p^2}).$$

Following the recipe in Section 6.4 we consider the following rational section of $\mathcal{O}_{C_{\mathbb{Z}/p^2}}(D_{0,3})$

$$l := \frac{10x^4 + x^3 + 17x + 14 + (15x + 9)y}{10x^4 + 16x^3 + 7x^2 + 7x + 10}.$$

since it generates $\mathcal{O}_{C_{\mathbb{Z}/p^2}}(D_{0,3})$ in a neighborhood of the supports of $\widetilde{G}_{t,2}$ and $E_{0,4}$. Then

$$\operatorname{div}(l) = 3 \cdot (-1, 1) + (17, 23) + (15, 10) - 2 \cdot (12, 23) - 2 \cdot (5, 20) - (0, 1) \quad \text{in } \operatorname{Div}(V_{1, \mathbb{Z}/p^2}) \subset \operatorname{Div}(C_{\mathbb{Z}/p^2}).$$

Hence by (6.4.4) the canonical isomorphism

$$\mathcal{M}(D_{0,3}, \widetilde{G}_{t,2}) = \widetilde{G}_{t,2}^* \mathcal{O}_{C_{\mathbb{Z}/p^2}}(D_{0,3}) \longrightarrow E_{0,4}^* \mathcal{O}_{C_{\mathbb{Z}/p^2}}(D_{0,3}) = \mathcal{M}(D_{0,3}, E_{0,4})$$

described in Section 6.4 is characterised by

$$(8.4.3) \quad \widetilde{G}_{t,2}^* l \longmapsto h_2(\operatorname{div}(l)) \cdot E_{0,4}^* l = 14 \cdot E_{0,4}^* l.$$

where

$$\begin{aligned} \widetilde{G}_{t,2}^* l &:= (\delta^* l)^{12} \otimes (\beta^* l)^{-6} \otimes (\infty^* l)^{-10} \otimes (\infty_-^* l)^4 = -(\delta^* 1)^{12} \otimes (\beta^* 1)^{-6} \otimes (\infty^* \frac{z^2}{z-3p})^{-10} \otimes (\infty_-^* 1)^4, \\ E_{0,4}^* l &:= (\beta^* l)^2 \otimes (\infty_4^* l) \otimes (\infty^* l)^{-1} \otimes (\infty_-^* l)^{-2} = 16 \cdot (\beta^* 1)^2 \otimes (\infty_4^* \frac{z^2}{z-3p}) \otimes (\infty^* \frac{z^2}{z-3p})^{-1} \otimes (\infty_-^* 1)^{-2}. \end{aligned}$$

Equations (8.4.2) and (8.4.3) imply that $\tilde{t} = -(1 + 2p) \cdot s_{D,E}((0, 3), (0, 4))$.

Let $\overline{A}_{\tilde{t}}, \overline{B}_{\tilde{t}}, \overline{C}$ and $\overline{D}_{\tilde{t}}$ be the compositions of the reduction map $\mathcal{M}^\times(\mathbb{Z}_p) \rightarrow \mathcal{M}(\mathbb{Z}/p^2)$ and respectively $A_{\tilde{t}}, B_{\tilde{t}}, C$ and $D_{\tilde{t}}$, defined in (4.2), (4.3) and (4.4). Using (6.6.13) and (8.4.1) we get, for n in \mathbb{Z}^2 ,

$$(8.4.4) \quad \begin{aligned} \overline{A}_{\tilde{t}}(n) &= (-1)^{n_2} (1 + (4n_2)t) \cdot s_{D,0}((0, 3), (2n_1, 4n_2)), \\ \overline{B}_{\tilde{t}}(n) &= (1 + (4n_2)p) s_{0,E}((4n_1, 3n_2), (0, 4)), \\ \overline{C}(n) &= (-1)^{n_2^2} (1 + (4n_1^2 + (4+4)n_1n_2 + 2n_2^2)p) \cdot s_{0,0}((4n_1, 3n_2), (2n_1, 4n_2)), \\ \overline{D}_{\tilde{t}}(n) &= -(1 + (4n_1^2 + 3n_1n_2 + 2n_2^2 + 3n_2 + 2)p) \cdot s_{D,E}((4n_1, 3 + 3n_2), (2n_1, 4 + 4n_2)), \\ \overline{\kappa}(n) &= -(1 + (4n_1^2 + 3n_1n_2 + 2n_2^2 + 2n_2 + 2)p) \cdot s_{D,E}((n_1, 3 + 2n_2), (3n_1, 4 + n_2)), \end{aligned}$$

hence, using the bijection (8.3.1),

$$(8.4.5) \quad \overline{\kappa}_1 = z_1, \quad \overline{\kappa}_2 = 3 + 2z_2, \quad \overline{\kappa}_3 = 4z_1^2 + 3z_1z_2 + 2z_2^2 + 2z_2 + 2.$$

8.5 The rational points with a specific image mod 5.

By 8.4.4 the image in $T(\mathbb{F}_p)$ of a point $\pm \overline{D}_{\tilde{t}}(n)$ for $n \in \mathbb{Z}^2$ is always of the form $\pm s_{D,E}(0, 0)$, hence, looking at (8.1.3) we see that there is no point $T(\mathbb{Z})$ with reduction $\widetilde{j}_{b,0}(\overline{\alpha}) \in T(\mathbb{F}_p)$. Hence $C(\mathbb{Z})_{\overline{\alpha}} = U_1(\mathbb{Z})_{\overline{\alpha}}$.

Let $f_1, f_2 \in \mathcal{O}(\widetilde{T}_t^p)^{\wedge p}$ be generators of the kernel of $\widetilde{j}_{b-1}^* : \mathcal{O}(\widetilde{T}_t^p)^{\wedge p} \rightarrow \mathcal{O}(\widetilde{U}_u^p)^{\wedge p}$ as in Section 4. The bijection (8.3.1) gives an isomorphism $\mathbb{F}_p \otimes \mathcal{O}(\widetilde{T}_t^p) = \mathbb{F}_p[\lambda_1, \lambda_2, \tau]$ and since the images $\overline{f_1}, \overline{f_2}$ of f_1, f_2 in $\mathbb{F}_p \otimes \mathcal{O}(\widetilde{T}_t^p)$ are generators of the kernel of $\widetilde{j}_{b,1}^* : \mathbb{F}_p \otimes \mathcal{O}(\widetilde{T}_t^p)^{\wedge p} \rightarrow \mathbb{F}_p \otimes \mathcal{O}(\widetilde{U}_u^p)^{\wedge p}$ we can suppose that

$$\overline{f_1} = \lambda_2, \quad \overline{f_2} = \tau - 1.$$

By (8.4.5) we have

$$\kappa^* \overline{f_1} = \overline{\kappa_2} = 3 + 2z_2, \quad \kappa^* \overline{f_2} = \overline{\kappa_3} - 1 = 4z_1^2 + 3z_1z_2 + 2z_2^2 + 2z_2 + 1.$$

Let A be $\mathbb{Z}_p\langle z_1, z_2 \rangle / (\kappa^* f_1, \kappa^* f_2)$. Then the ring

$$(8.5.1) \quad \overline{A} := A/pA = \mathbb{F}_p[z_1, z_2] / (\kappa^* \overline{f_1}, \kappa^* \overline{f_2}) = \mathbb{F}_p[z_1, z_2] / (z_2 - 1, 4z_1^2 + 3z_1)$$

has dimension 2 over \mathbb{F}_p , hence by Theorem 4.12 $U(\mathbb{Z})_{\overline{\alpha}}$ contains at most 2 points. Since both

$$\alpha \quad \text{and} \quad (12/7, 20/7) \in V_1(\mathbb{Z}[1/7])$$

reduce to $\overline{\alpha}$ we deduce that $C(\mathbb{Z})_{\overline{\alpha}} = U_1(\mathbb{Z})_{\overline{\alpha}}$ is made of the these two points.

8.6 Determination of all rational points

Denoting $(3, -1) \in V_1(\mathbb{F}_p) \subset C(\mathbb{F}_p)$ as ε we have

$$C(\mathbb{F}_p) = \{\overline{\infty}, \overline{\infty}_-, \overline{\alpha}, \iota(\overline{\alpha}), \eta(\overline{\alpha}), (\iota \circ \eta)(\overline{\alpha}), \overline{\gamma}, \iota(\overline{\gamma}), \eta(\overline{\gamma}), (\iota \circ \eta)(\overline{\gamma}), \varepsilon, \iota(\varepsilon)\}.$$

Using that for any point Q in $C(\mathbb{F}_p)$ the condition $T(\mathbb{Z})_{\widetilde{j}_{b,i}(Q)} = \emptyset$ implies $U_i(\mathbb{Z})_Q = \emptyset$ we get

$$U_0(\mathbb{Z})_{\overline{\infty}} = U_0(\mathbb{Z})_{\overline{\infty}_-} = U_1(\mathbb{Z})_{\varepsilon} = U_1(\mathbb{Z})_{\iota(\varepsilon)} = U_1(\mathbb{Z})_{\overline{\gamma}} = U_1(\mathbb{Z})_{\eta(\overline{\gamma})} = U_1(\mathbb{Z})_{\iota(\eta(\overline{\gamma}))} = U_1(\mathbb{Z})_{\eta(\eta(\overline{\gamma}))} = \emptyset.$$

Applying our method to $\overline{\infty}$ we discover that $U_1(\mathbb{Z})_{\overline{\infty}}$ contains at most 2 points and the same holds for $U_1(\mathbb{Z})_{\overline{\infty}_-}$. Moreover the action of $\langle \eta, \iota \rangle$ on $C(\mathbb{Z})$ tells that $U_1(\mathbb{Z})_{\iota(\overline{\alpha})}$, $U_1(\mathbb{Z})_{\eta(\overline{\alpha})}$ and $U_1(\mathbb{Z})_{\eta(\iota(\overline{\alpha}))}$ are sets containing exactly 2 elements. Hence

$$U_1(\mathbb{Z}) = U_1(\mathbb{Z})_{\overline{\alpha}} \cup U_1(\mathbb{Z})_{\iota(\overline{\alpha})} \cup U_1(\mathbb{Z})_{\eta(\overline{\alpha})} \cup U_1(\mathbb{Z})_{\eta(\iota(\overline{\alpha}))} \cup U_1(\mathbb{Z})_{\overline{\infty}_-} \cup U_1(\mathbb{Z})_{\overline{\infty}}$$

contains at most 12 elements. Looking at the orbits of the action of $\langle \eta, \iota \rangle$ on $U_1(\mathbb{Z})$ we see that $\#U_1(\mathbb{Z}) \equiv 2 \pmod{4}$, hence $\#U_1(\mathbb{Z}) \leq 10$. Since $U_1(\mathbb{Z})$ contains $\overline{\infty}, \overline{\infty}_-$ and all the images by $\langle \eta, \iota \rangle$ of $U_1(\mathbb{Z})_{\overline{\alpha}}$ we conclude that $\#U_1(\mathbb{Z}) = 10$.

Applying our method to the point $\overline{\gamma}$ we see that $U_0(\mathbb{Z})_{\overline{\gamma}}$ contains at most two points, one of them being γ . Moreover solving the equations $\kappa^* \overline{f_i} = 0$ we see that if there is another point γ' in $U_0(\mathbb{Z})_{\overline{\gamma}}$ then there exist $n_1, n_2 \in \mathbb{Z}$ such that

$$j_b(\gamma') = 39G_1 + 17G_2 + 5n_1\widetilde{G}_1 + 5n_2\widetilde{G}_2.$$

Confronting this information with the image of the map $j_b : C(\mathbb{F}_7) \rightarrow J(\mathbb{F}_7)$ we derive a contradiction, hence $U_0(\mathbb{Z})_{\overline{\gamma}} = \{\gamma\}$. Applying our method to ε we see that $U_0(\mathbb{Z})_{\varepsilon}$ contains at most 2 points corresponding to two different solutions to the equations $\kappa^* \overline{f_i} = 0$. We can see

that one of the two solutions does not lift to a point in $U_0(\mathbb{Z})_\varepsilon$ in the same way we excluded the existence of $\gamma' \in U_0(\mathbb{Z})_{\bar{\gamma}}$. Hence $U_0(\mathbb{Z})_\varepsilon$ has cardinality at most 1. Using that for every $Q \in C(\mathbb{F}_p)$ and every automorphism ω of C we have $\#U_0(\mathbb{Z})_Q = \#U_0(\mathbb{Z})_{\omega(Q)}$, we deduce that

$$U_0(\mathbb{Z}) = U_0(\mathbb{Z})_{\bar{\gamma}} \cup U_0(\mathbb{Z})_{\iota(\bar{\gamma})} \cup U_0(\mathbb{Z})_{\eta(\bar{\gamma})} \cup U_0(\mathbb{Z})_{\eta\iota(\bar{\gamma})} \cup U_0(\mathbb{Z})_\varepsilon \cup U_0(\mathbb{Z})_{\iota(\varepsilon)}$$

contains at most 6 points. Looking at the orbits of the action of $\langle \eta, \iota \rangle$ on $U_0(\mathbb{Z})$ we see that $\#U_0(\mathbb{Z}) \equiv 0 \pmod{4}$, hence $\#U_4(\mathbb{Z}) \leq 4$, and since $U_0(\mathbb{Z})$ contains the orbit of γ we conclude that $\#U_0(\mathbb{Z}) = 4$. Finally

$$\#C(\mathbb{Z}) = \#U_0(\mathbb{Z}) + \#U_1(\mathbb{Z}) = 4 + 10 = 14.$$

9 Some further remarks

This section was added later, because a referee wanted to know about §9.2 and §9.4.

9.1 Complex uniformisations of some of the objects involved

Let C be a projective curve over \mathbb{Q} , smooth, and geometrically irreducible, and let g be its genus. The universal cover of $P^\times(\mathbb{C})$ is described in [6], Propositions 4.5 and 4.6. The covering space, denoted D_τ , is $M_{1,g}(\mathbb{C}) \times M_{g,1}(\mathbb{C}) \times \mathbb{C}$, hence a \mathbb{C} -vector space of dimension $2g + 1$. The biextension structure on $M_{1,g}(\mathbb{C}) \times M_{g,1}(\mathbb{C}) \times \mathbb{C}$ is trivial, that is, for all x, x_1, x_2 in $M_{1,g}(\mathbb{C})$, all y, y_1, y_2 in $M_{g,1}(\mathbb{C})$, and all z_1, z_2 in \mathbb{C} , we have:

$$(9.1.1) \quad \begin{aligned} (x_1, y, z_1) +_1 (x_2, y, z_2) &= (x_1 + x_2, y, z_1 + z_2), \\ (x, y_1, z_1) +_2 (x, y_2, z_2) &= (x, y_1 + y_2, z_1 + z_2). \end{aligned}$$

The fundamental group $\pi_1(P^\times(\mathbb{C}), 1)$ is

$$(9.1.2) \quad Q^u(\mathbb{Z}) := \left\{ \begin{pmatrix} 1 & x & z \\ 0 & 1_{2g} & y \\ 0 & 0 & 1 \end{pmatrix} : x \in M_{1,2g}(\mathbb{Z}), y \in M_{2g,1}(\mathbb{Z}), z \in \mathbb{Z} \right\},$$

also known as a Heisenberg group. Its action on D_τ is given in [6, (4.5.3)].

Now recall the definition of T in (2.12). As $M_{2g,1}(\mathbb{Z})$ is the lattice of $J(\mathbb{C})$, and $M_{1,2g}(\mathbb{Z})$ the lattice of $J^\vee(\mathbb{C})$, each f_i is given by an antisymmetric matrix $f_{i,\mathbb{Z}}$ in $M_{2g,2g}(\mathbb{Z})$ such that for all y in $M_{2g,1}(\mathbb{Z})$ we have $f_i(y) = y^t \cdot f_{i,\mathbb{Z}}$, and by a complex matrix $f_{i,\mathbb{C}}$ in $M_{g,g}(\mathbb{C})$ such that for all v in $M_{g,1}(\mathbb{C})$, for each i we have $f_i(v) = v^t \cdot f_{i,\mathbb{C}}$ in $M_{1,g}(\mathbb{C})$. For more details about this description of the f_i see the beginning of [6, §4.7]. Then we have

$$(9.1.3) \quad \pi_1(T(\mathbb{C})) = \left\{ \begin{pmatrix} 1_{\rho-1} & m \cdot f(y) & z \\ 0 & 1_{2g} & y \\ 0 & 0 & 1 \end{pmatrix} : y \in M_{2g,1}(\mathbb{Z}), z \in M_{\rho-1,1}(\mathbb{Z}) \right\},$$

with $m \cdot f(y) \in M_{\rho-1,2g}(\mathbb{Z})$ with rows the $m \cdot y^t \cdot f_{i,\mathbb{Z}}$. So, $\pi_1(T(\mathbb{C}))$ is a central extension of $M_{2g,1}(\mathbb{Z})$ by $M_{\rho-1,1}(\mathbb{Z})$, with commutator pairing sending (y, y') to $(2my^t \cdot f_{i,\mathbb{Z}} \cdot y')_i$.

The universal covering $\widetilde{T(\mathbb{C})}$ is given by

$$(9.1.4) \quad \begin{aligned} \widetilde{T(\mathbb{C})} &= \{(m \cdot (c + f(v)), v, w) : v \in M_{g,1}(\mathbb{C}), w \in M_{\rho-1,1}(\mathbb{C})\} \\ &\subset M_{\rho-1,g}(\mathbb{C}) \times M_{1,g}(\mathbb{C}) \times M_{\rho-1,1}(\mathbb{C}), \end{aligned}$$

with $m \cdot (c + f(v)) \in M_{\rho-1,g}(\mathbb{C})$ with rows the $m \cdot (\tilde{c}_i + v^t \cdot f_{i,\mathbb{C}})$ with \tilde{c}_i a lift of c_i in $M_{1,g}(\mathbb{C})$. The action of $\pi_1(T(\mathbb{C}), 1)$ on $\widetilde{T(\mathbb{C})}$ is given again, with the necessary changes, by [6, (4.5.3)].

Now that we know $\pi_1(T(\mathbb{C}), 1)$ we investigate which quotient of $\pi_1(C(\mathbb{C}), b)$ it is, via $\tilde{j}_b: C(\mathbb{C}) \rightarrow T(\mathbb{C})$. We consider the long exact sequence of homotopy groups induced by the $\mathbb{C}^{\times, \rho-1}$ -torsor $T(\mathbb{C}) \rightarrow J(\mathbb{C})$, taking into account that $\mathbb{C}^{\times, \rho-1}$ is connected and that $\pi_2(J(\mathbb{C})) = 0$:

$$(9.1.5) \quad \pi_1(\mathbb{C}^{\times, \rho-1}, 1) \hookrightarrow \pi_1(T(\mathbb{C}), 1) \twoheadrightarrow \pi_1(J(\mathbb{C}), 0).$$

Again we see that $\pi_1(T(\mathbb{C}), 1)$ is a central extension of the free abelian group $\pi_1(J(\mathbb{C}), 0)$ by $\mathbb{Z}^{\rho-1}$, and from the matrix description we know that the i th coordinate of the commutator pairing is given by $m f_i: H_1(J(\mathbb{C}), \mathbb{Z}) \rightarrow H_1(J^\vee(\mathbb{C}), \mathbb{Z}) = H_1(J(\mathbb{C}), \mathbb{Z})^\vee$. The \mathbb{Z} -module of antisymmetric \mathbb{Z} -valued pairings on $H_1(J^\vee(\mathbb{C}), \mathbb{Z})$ is $\bigwedge^2 H^1(J(\mathbb{C}), \mathbb{Z}) = H^2(J(\mathbb{C}), \mathbb{Z})$, and $m f_i$ is the cohomology class (first Chern class) of the \mathbb{C}^\times -torsor T_i :

$$(9.1.6) \quad m f_i = c_1(T_i) \quad \text{in } H^2(J(\mathbb{C}), \mathbb{Z}).$$

There is a central extension

$$(9.1.7) \quad H_2(J(\mathbb{C}), \mathbb{Z}) \hookrightarrow E \twoheadrightarrow \pi_1(J(\mathbb{C}), 0)$$

that is universal in the sense that every central extension of $\pi_1(J(\mathbb{C}), 0)$ by a free abelian group arises by pushout from $H_2(J(\mathbb{C}), \mathbb{Z})$. We denote

$$(9.1.8) \quad G := \pi_1(C(\mathbb{C}), b).$$

The map $j_b: C \rightarrow J$ gives $G \rightarrow \pi_1(J(\mathbb{C}), 0)$, and this is the maximal abelian quotient. The second quotient in the descending central series of G gives the central extension:

$$(9.1.9) \quad [G, G]/[G, [G, G]] \hookrightarrow G/[G, [G, G]] \twoheadrightarrow G/[G, G] = G^{\text{ab}} = \pi_1(J(\mathbb{C}), 0).$$

This extension (9.1.9) arises from (9.1.7) by pushout via a morphism from $H_2(J(\mathbb{C}), \mathbb{Z})$ to $[G, G]/[G, [G, G]]$:

$$(9.1.10) \quad \begin{array}{ccccc} H_2(J(\mathbb{C}), \mathbb{Z}) & \hookrightarrow & E & \twoheadrightarrow & G^{\text{ab}} \\ \downarrow & & \downarrow & & \parallel \\ [G, G]/[G, [G, G]] & \hookrightarrow & G/[G, [G, G]] & \twoheadrightarrow & G^{\text{ab}}. \end{array}$$

The left vertical arrow is surjective because commutators of lifts in E of elements of G^{ab} are mapped to the commutators of lifts in $G/[G, [G, G]]$, and so give generators of $[G, G]/[G, [G, G]]$.

From the usual presentation of G with generators $\alpha_1, \beta_1, \dots, \alpha_g, \beta_g$, with the only relation $[\alpha_1, \beta_1] \cdots [\alpha_g, \beta_g] = 1$, we see that the obstruction in lifting $G \rightarrow G^{\text{ab}}$ to $G \rightarrow E$ in the top row

of (9.1.10) is the image of $[\alpha_1, \beta_1] \cdots [\alpha_g, \beta_g]$ in $H_2(J(\mathbb{C}), \mathbb{Z})$. This image is a generator of the image of $H_2(C(\mathbb{C}), \mathbb{Z})$ under j_b . So the pushout in (9.1.10) factors through the pushout by the quotient of $H_2(J(\mathbb{C}), \mathbb{Z})$ by $H_2(C(\mathbb{C}), \mathbb{Z})$:

$$(9.1.11) \quad \begin{array}{ccccc} H_2(J(\mathbb{C}), \mathbb{Z})/H_2(C(\mathbb{C}), \mathbb{Z}) & \hookrightarrow & E' & \twoheadrightarrow & G^{\text{ab}} \\ \downarrow & & \downarrow & & \parallel \\ [G, G]/[G, [G, G]] & \hookrightarrow & G/[G, [G, G]] & \twoheadrightarrow & G^{\text{ab}}. \end{array}$$

Using again the presentation of G we can split this morphism of extensions, and, using that $H_2(J(\mathbb{C}), \mathbb{Z})/H_2(C(\mathbb{C}), \mathbb{Z})$ is generated by commutators of lifts of elements of G^{ab} , conclude that all vertical arrows in (9.1.11) are isomorphisms.

In particular, we have that $[G, G]/[G, [G, G]]$ is the same as $H_2(J(\mathbb{C}), \mathbb{Z})/H_2(C(\mathbb{C}), \mathbb{Z})$. From (9.1.6) we see that the sub- \mathbb{Z} -module of $H^2(J(\mathbb{C}), \mathbb{Z}(1))$ (note the Tate twist, now we take the Hodge structures into account) spanned by the mf_i is obtained in 4 steps: take the kernel of $H^2(J(\mathbb{C}), \mathbb{Z}(1)) \rightarrow H^2(C(\mathbb{C}), \mathbb{Z}(1))$, take the $(0, 0)$ -part, then $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ acts, through the Galois group of a finite extension of \mathbb{Q} , take the invariants, then take the image of the multiplication by m on that.

Dually, this means that $\pi_1(T(\mathbb{C}), 1)$ arises as the pushout

$$(9.1.12) \quad \begin{array}{ccccc} H_2(J(\mathbb{C}), \mathbb{Z}(-1))/H_2(C(\mathbb{C}), \mathbb{Z}(-1)) & \hookrightarrow & G/[G, [G, G]] & \twoheadrightarrow & G^{\text{ab}} \\ \downarrow & & \downarrow & & \parallel \\ ((H_2(J(\mathbb{C}), \mathbb{Z}(-1))/H_2(C(\mathbb{C}), \mathbb{Z}(-1)))_{(0,0)})_{\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})} & \hookrightarrow & \pi_1(T(\mathbb{C}), 1) & \twoheadrightarrow & G^{\text{ab}}, \end{array}$$

where the subscript $(0, 0)$ means the largest quotient of type $(0, 0)$, where the subscript $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ means co-invariants modulo torsion, and where the left vertical map is m times the quotient map. We repeat that the morphism from $\pi_1(C(\mathbb{C})) = G$ to $\pi_1(T(\mathbb{C}), 1)$ given by the middle vertical map is induced by $\tilde{j}_b: C(\mathbb{C}) \rightarrow J(\mathbb{C})$.

9.2 Finiteness of rational points

In this section we reprove Faltings's finiteness result [13] in the special case where $r < g + \rho - 1$. This was already done in [3], Lemma 3.2 (where the base field is either \mathbb{Q} or imaginary quadratic). We begin by collecting some ingredients on good formal coordinates of the \mathbb{G}_m -biextension $P^{\times, \rho-1} \rightarrow J \times J^{\vee, \rho-1}$ over \mathbb{Q} , and on what C looks like in such coordinates.

9.2.1 Formal trivialisations

Let A, B and G be connected smooth commutative group schemes over a field $k \supset \mathbb{Q}$, and let $E \rightarrow A \times B$ be a commutative G -biextension. Let a be in $A(k)$, $b \in B(k)$ and $e \in E(k)$. For $n \in \mathbb{N}$, let $A^{a,n}$ be the n th infinitesimal neighborhood of a in A , hence its coordinate ring is $\mathcal{O}_{A,a}/m_a^{n+1}$. We use similar notation for B with b , and E with e , and also for the points 0 of A, B and E , and, similarly, the formal completion of A at a is denoted by $A^{a,\infty}$, etc. We also use such notation in a relative context, for example, for the group schemes $E \rightarrow B$

and $E \rightarrow A$. We view completions as $A^{a,\infty}$ as set-valued functors on the category of local k -algebras with residue field k such that every element of the maximal ideal is nilpotent. For such a k -algebra R , $A^{a,\infty}(R)$ is the inverse image of a under $A(R) \rightarrow A(k)$. Then $A^{0,\infty}$ is the formal group of A .

We now want to show that the formal $G^{0,\infty}$ -biextension $E^{0,\infty} \rightarrow A^{0,\infty} \times B^{0,\infty}$ is isomorphic to the trivial biextension (the object $G^{0,\infty} \times A^{0,\infty} \times B^{0,\infty}$ with $+_1$ given by addition on the 1st and 2nd coordinate, and $+_2$ by addition on the 1st and 3rd coordinate). As exp for $A^{0,\infty}$ gives a functorial isomorphism $T_{A/k}(0) \otimes_k \mathbb{G}_{a,k}^{0,\infty} \rightarrow A^{0,\infty}$, and similarly for B and G , it suffices to prove this triviality for $\mathbb{G}_a^{0,\infty}$ -biextensions of $\mathbb{G}_a^{0,\infty} \times \mathbb{G}_a^{0,\infty}$ over k . One easily checks that the group of automorphisms of the trivial $\mathbb{G}_a^{0,\infty}$ -biextension of $\mathbb{G}_a^{0,\infty} \times \mathbb{G}_a^{0,\infty}$ over k that induce the identity on all three $\mathbb{G}_a^{0,\infty}$'s is $(k, +)$, with $c \in k$ acting as $(g, a, b) \mapsto (g + cab, a, b)$. As this group is commutative, it then follows that the group of automorphisms of the $G^{0,\infty}$ -biextension $E^{0,\infty} \rightarrow A^{0,\infty} \times B^{0,\infty}$ that induce identity on $G^{0,\infty}$, $A^{0,\infty}$, and $B^{0,\infty}$, is equal to the k -vector space of k -bilinear maps $T_{A/k}(0) \times T_{B/k}(0) \rightarrow T_{G/k}(0)$. This indicates how to trivialise $E^{0,\infty}$. We choose a section \tilde{e} of the G -torsor $E \rightarrow A \times B$ over the closed subscheme $A^{0,1} \times B^{0,1}$ of $A \times B$:

$$\begin{array}{ccc} & & E \\ & \nearrow \tilde{e} & \downarrow \\ A^{0,1} \times B^{0,1} & \longrightarrow & A \times B, \end{array} \quad \text{with } \tilde{e}(0,0) = e \text{ in } E(k).$$

Note that

$$\mathcal{O}(A^{0,1} \times B^{0,1}) = (k \oplus m_{A^{0,1}}) \otimes (k \oplus m_{B^{0,1}}) = k \oplus m_{A^{0,1}} \oplus m_{B^{0,1}} \oplus (m_{A^{0,1}} \otimes m_{B^{0,1}}).$$

Hence two such \tilde{e} differ by a k -algebra morphism from $k \oplus m_{G^{0,2}} = k \oplus m_{G^{0,1}} \oplus \text{Sym}^2 m_{G^{0,1}}$ (use the exponential map) to $k \oplus m_{A^{0,1}} \oplus m_{B^{0,1}} \oplus (m_{A^{0,1}} \otimes m_{B^{0,1}})$, hence by a triple of k -linear maps from $m_{G^{0,1}}$ to $m_{A^{0,1}}$, $m_{B^{0,1}}$, and $m_{A^{0,1}} \otimes m_{B^{0,1}}$. The linear maps $m_{G^{0,1}} \rightarrow m_{A^{0,1}}$ and $m_{G^{0,1}} \rightarrow m_{B^{0,1}}$ correspond to the differences on $A^{0,1} \times B^{0,0}$ and on $A^{0,0} \times B^{0,1}$, respectively. There are unique such linear maps such that the adjusted \tilde{e} is compatible with the given trivialisations of $E \rightarrow A \times B$ over $A^{0,1} \times B^{0,0}$ and over $A^{0,0} \times B^{0,1}$. In geometric terms, \tilde{e} , assumed to be adjusted, is then a splitting of $T_G(0)_B \hookrightarrow T_{E/B}(0) \twoheadrightarrow T_A(0)_B$ over $B^{0,1}$ that is compatible with the already given splitting over $0 \in B(k)$, and it is also a splitting of $T_G(0)_A \hookrightarrow T_{E/A}(0) \twoheadrightarrow T_B(0)_A$ over $A^{0,1}$ that is compatible with the already given splitting over $0 \in A(k)$. The splitting over $B^{0,1}$ gives an isomorphism from $(T_G(0) \oplus T_A(0))_{B^{0,1}}$ to $(T_{E/B})_{B^{0,1}}$. So the exponential map, for $+_1$, for the pullback to $B^{0,1}$ of $E \rightarrow B$, gives an isomorphism of formal groups over $B^{0,1}$:

$$((T_G(0) \oplus T_A(0)) \otimes_k \mathbb{G}_a^{0,\infty})_{B^{0,1}} \hookrightarrow E_{B^{0,1}}^{0,\infty}.$$

Viewing $E_{B^{0,1}}^{0,\infty}$ as the tangent space at the zero section of the pullback to $A^{0,\infty}$ of $E \rightarrow A$, this isomorphism gives a splitting of $T_G(0)_A \hookrightarrow T_{E/A}(0) \twoheadrightarrow T_B(0)_A$ over $A^{0,\infty}$. The exponential map for $+_2$ for the pullback to $A^{0,\infty}$ of $E \rightarrow A$ then gives an isomorphism of formal groups over $A^{0,\infty}$:

$$G^{0,\infty} \times B^{0,\infty} \times A^{0,\infty} \xlongequal{\quad} (G^{0,\infty} \times B^{0,\infty})_{A^{0,\infty}} \hookrightarrow E_{A^{0,\infty}/A^{0,\infty}}^{0,\infty} \xlongequal{\quad} E^{0,\infty},$$

where $E_{A^{0,\infty}/A^{0,\infty}}^{0,\infty}$ denotes the completion along the zero section of the pullback via $A^{0,\infty} \rightarrow A$ of $E \rightarrow A$. The compatibility between $+_1$ and $+_2$ on E ensures that this isomorphism is an isomorphism of biextensions, with the trivial biextension structure on the left.

Now that we know what good formal coordinates at 0 in $E(k)$ are, we look at the point e in $E(k)$, over (a, b) in $(A \times B)(k)$. We produce an isomorphism $E^{0,\infty} \rightarrow E^{e,\infty}$, using the partial group laws. Let E_b be the fibre over b of $E \rightarrow B$. We choose a section

$$\begin{array}{ccc} & & E_b \\ & \nearrow \tilde{e}_1 & \downarrow \\ A^{a,1} \times \{b\} & \longrightarrow & A \times \{b\} \end{array} \quad \text{with } \tilde{e}_1(a, b) = e \text{ in } E(k).$$

The exponentials for the group laws of E_b and A then give a section

$$\begin{array}{ccc} & & E_b \\ & \nearrow \tilde{e}_1^\infty & \downarrow \\ A^{a,\infty} \times \{b\} & \longrightarrow & A \times \{b\}, \end{array}$$

that we view as an $A^{a,\infty}$ -valued point of E_b , and as a section of the group scheme $E_{A^{a,\infty}} \rightarrow A^{a,\infty}$, with group law $+_2$. The translation by \tilde{e}_1^∞ on this group scheme induces translation by b on $B_{A^{a,\infty}}$, and maps $(a, 0)$, the 0 element of E_a , to e . Hence it induces an isomorphism of formal schemes $E^{(a,0),\infty} \rightarrow E^{e,\infty}$. In order to get an isomorphism $E^{0,\infty} \rightarrow E^{(a,0),\infty}$, we repeat the process above, but with the roles of A and B exchanged. We choose a section $\tilde{\theta}_2: \{a\} \times B^{0,1} \rightarrow E_a$ of $E_a \rightarrow \{a\} \times B$. Then the exponential for $+_2$ gives us a section $\tilde{\theta}_2^\infty: \{a\} \times B^{0,\infty} \rightarrow E_a$ of $E_a \rightarrow \{a\} \times B$. This $\tilde{\theta}_2^\infty$ is a section of the group scheme $E_{B^{0,\infty}} \rightarrow B^{0,\infty}$, and the translation on it by $\tilde{\theta}_2^\infty$ sends 0 in $E(k)$ to $(a, 0)$, hence gives an isomorphism of formal schemes $E^{0,\infty} \rightarrow E^{(a,0),\infty}$. Composition then gives us an isomorphism $E^{0,\infty} \rightarrow E^{e,\infty}$, and the good formal coordinates on E at $0 \in E(k)$ give what we call good formal coordinates at e . Similarly, we get a section $\tilde{\theta}_1^\infty$ of $E_{A^{0,\infty}} \rightarrow A^{0,\infty}$ and a section \tilde{e}_2^∞ of $E_{B^{b,\infty}} \rightarrow B^{b,\infty}$ giving isomorphisms $E^{0,\infty} \rightarrow E^{(0,b),\infty}$ and $E^{(0,b),\infty} \rightarrow E^{e,\infty}$, hence by composition a 2nd isomorphism $E^{0,\infty} \rightarrow E^{e,\infty}$. These isomorphisms are equal for a unique choice of $\tilde{\theta}_1$ and \tilde{e}_2 (given the choices of $\tilde{\theta}_2$ and \tilde{e}_1).

In Section 9.2.3 we will use that these isomorphisms transport all additions that occur in (4.4) to additions in $E^{0,\infty}$ and therefore to additions in the trivial formal biextension.

9.2.2 Zariski density of the curve in formally trivial coordinates

Let C be as in the beginning of Section 2. Let $\widetilde{C}(\mathbb{C})$ be the inverse image of $C(\mathbb{C})$ under the universal cover $\widetilde{T}(\mathbb{C}) \rightarrow T(\mathbb{C})$. Then $\widetilde{C}(\mathbb{C})$ is connected since $\tilde{j}_b: C \rightarrow T$ gives a surjection on complex fundamental groups. Now we consider the complex analytic variety $\widetilde{T}(\mathbb{C})$ as a complex algebraic variety via the bijection $\widetilde{T}(\mathbb{C}) = \mathbb{C}^{g+\rho-1}$ as given in (9.1.4). The analytic subset $\widetilde{C}(\mathbb{C})$ contains the orbit of 0 under $\pi_1(T(\mathbb{C}), 1)$. This orbit surjects to the lattice of $J(\mathbb{C})$ in $M_{g,1}(\mathbb{C})$, and over each lattice point, its fibre in $M_{\rho-1,1}(\mathbb{C})$ is contains a translate of $2\pi i M_{\rho-1,1}(\mathbb{Z})$. Hence this orbit is Zariski dense in $\mathbb{C}^{g+\rho-1}$. It follows that the formal completion of $\widetilde{C}(\mathbb{C})$ at any of its points is Zariski dense in $\mathbb{C}^{g+\rho-1}$: if a polynomial function on $\mathbb{C}^{g+\rho-1}$ is zero on such a completion, then it vanishes on the connected component of $\widetilde{C}(\mathbb{C})$ of that point, hence on $\widetilde{C}(\mathbb{C})$.

We express our conclusion in more algebraic terms: for $c \in C(\mathbb{C})$, with images $t \in T(\mathbb{C})$ and in $P^{\times, \rho-1}(\mathbb{C})$, each polynomial in good formal coordinates at t of the biextension $P^{\times, \rho-1} \rightarrow J \times J^\vee$ over \mathbb{C} that vanishes on $\tilde{j}_b(C_{\mathbb{C}}^{e, \infty})$, vanishes on $T_{\mathbb{C}}^{t, \infty}$. This statement then also holds with \mathbb{C} replaced by any subfield, or even any subring of the form $\mathbb{Z}_{(p)}$ with p a prime number, or the localisation of $\overline{\mathbb{Z}}$ (the integral closure of \mathbb{Z} in \mathbb{C}) at a maximal ideal.

9.2.3 The p -adic closure in good formal coordinates

We stay in the situation of Section 2, but we denote $G := \mathbb{G}_m^{\rho-1}$, $A := J$ and $B := J^{\vee, 0\rho-1}$, and $E := P^{\times, \rho-1}$. Let d_G , d_A , and d_B be their dimensions: $d_G = \rho - 1$, $d_A = g$ and $d_B = (\rho - 1)g$.

Let $p > 2$ be a prime number. From Section 9.2.1 and Lemma 5.1.1 we conclude that we can choose *formal* parameters for E at 0, over $\mathbb{Z}_{(p)}$, such that they converge on the residue polydisk $E(\mathbb{Z}_p)_{\overline{0}}$, and such that they induce the trivial biextension structure on $\mathbb{Z}_p^{d_G} \times \mathbb{Z}_p^{d_A} \times \mathbb{Z}_p^{d_B}$. We keep the notation of Section 9.2.1, for e in $E(\mathbb{Z}_p)$, lying over (a, b) in $(A \times B)(\mathbb{Z}_p)$. This e plays the role that \tilde{t} has at the beginning of Section 4. As explained at the end of Section 9.2.1, we may and do assume that e is in $E(\mathbb{Z}_p)_{\overline{0}}$, and hence $a \in A(\mathbb{Z}_p)_{\overline{0}}$ and $b \in B(\mathbb{Z}_p)_{\overline{0}}$.

Assume now that, as in Section 4, for $i, j \in \{1, \dots, r\}$, we have x_i in $A(\mathbb{Z}_p)_{\overline{0}}$ and y_j in $B(\mathbb{Z}_p)_{\overline{0}}$, and $e_{i,j}$ in $E(\mathbb{Z}_p)_{\overline{0}}$ over (x_i, y_j) , and r_i in $E(\mathbb{Z}_p)_{\overline{0}}$ over (x_i, b) and s_j in $E(\mathbb{Z}_p)_{\overline{0}}$ over (a, y_j) . We denote the images under of all these elements under the bijection

$$E(\mathbb{Z}_p)_{\overline{0}} \longrightarrow \mathbb{Z}_p^{d_G} \times \mathbb{Z}_p^{d_A} \times \mathbb{Z}_p^{d_B}$$

as follows:

$$\begin{aligned} x_i &\mapsto (0, x_i, 0), & y_j &\mapsto (0, 0, y_j), & e_{i,j} &\mapsto (g_{i,j}, x_i, y_j) \\ r_i &\mapsto (r'_i, x_i, b), & s_j &\mapsto (s'_j, a, y_j), & e &\mapsto (e', a, b). \end{aligned}$$

Then a straightforward computation shows that the image of $D(n)$ as defined in (4.4) is

$$\left(e' + \sum_i n_i r'_i + \sum_j n_j s'_j + \sum_{i,j} n_i n_j g_{i,j}, a + \sum_i n_i x_i, b + \sum_j n_j y_j \right) \text{ in } \mathbb{Z}_p^{d_G} \times \mathbb{Z}_p^{d_A} \times \mathbb{Z}_p^{d_B}.$$

The conclusion is that in these coordinates, the map

$$\kappa: \mathbb{Z}_p^r \longrightarrow \mathbb{Z}_p^{d_G} \times \mathbb{Z}_p^{d_A} \times \mathbb{Z}_p^{d_B}$$

is a polynomial map, hence the Zariski closure of its image is an algebraic variety of dimension at most r .

9.2.4 Proof of finiteness

The proof is by contradiction. So assume that $r < g + \rho - 1$, and that $C(\mathbb{Q})$ is infinite. Let $p > 2$ be a prime number. Then there is a $u \in C(\mathbb{F}_p)$ such that the residue disk $C(\mathbb{Z}_p)_u$ contains infinitely many elements of $C(\mathbb{Q})$, hence infinitely many elements in the image of κ of Section 4.10. By construction, $\kappa(\mathbb{Z}_p^r)$ is contained in $T(\mathbb{Z}_p)_t$. The image of $T(\mathbb{Z}_p)_t$ in $\mathbb{Z}_p^{d_G} \times \mathbb{Z}_p^{d_A} \times \mathbb{Z}_p^{d_B}$ is $\mathbb{Z}_p^{\rho-1} \times \mathbb{Z}_p^g$, with \mathbb{Z}_p^g embedded in $\mathbb{Z}_p^{d_A} \times \mathbb{Z}_p^{d_B}$ as a sub- \mathbb{Z}_p -module. By the previous section, the

Zariski closure of $\kappa(\mathbb{Z}_p^r)$ in $\mathbb{Z}_p^{d_G} \times \mathbb{Z}_p^{d_A} \times \mathbb{Z}_p^{d_B}$ is of dimension at most r . Hence there are non-zero polynomial functions on $\mathbb{Z}_p^{p-1} \times \mathbb{Z}_p^g$ that are zero on infinitely many points of $C(\mathbb{Z}_p)_u$, and hence are zero on a non-empty open smaller disk. This contradicts, via a ring morphism $\mathbb{Z}_p \rightarrow \mathbb{C}$, the conclusion of Section 9.2.2.

9.3 The relation with p -adic heights

Let $p > 2$ be a prime number of good reduction for C . We consider the Poincaré torsor as \mathcal{M}^\times on $(J \times J)_{\mathbb{Q}_p}$ via (6.3.3), and we use the description of \mathcal{M}^\times given in (6.3.13).

Let \mathcal{D} be the subset $\text{Div}^0(C_{\mathbb{Q}_p}) \times \text{Div}^0(C_{\mathbb{Q}_p})$ made of pairs of divisors (D_1, D_2) having disjoint support. Let W be an isotropic complement of $\Omega_{C_{\mathbb{Q}_p}/\mathbb{Q}_p}^1(C_{\mathbb{Q}_p})$ in $H_{\text{dR}}^1(C_{\mathbb{Q}_p}/\mathbb{Q}_p)$ and let $\log: \mathbb{Q}_p^\times \rightarrow \mathbb{Q}_p$ be a group morphism extending the formal logarithm on $1 + p\mathbb{Z}_p$. With these choices made, Coleman and Gross ([9, (5.1)]) define the function (there denoted $\langle \cdot, \cdot \rangle$)

$$h_p: \mathcal{D} \rightarrow \mathbb{Q}_p,$$

the p -part of the p -adic height pairing. We define the function

$$\psi: \mathcal{M}^\times(\mathbb{Q}_p) \longrightarrow \mathbb{Q}_p$$

by demanding that for every effective D_1 and D_2 in $\text{Div}(C_{\mathbb{Q}_p})$ of the same degree and every E in $\text{Div}^0(C_{\mathbb{Q}_p})$, and every λ in \mathbb{Q}_p^\times , the element

$$\lambda \cdot \text{Norm}_{D_1/\mathbb{Q}_p}(1) \otimes \text{Norm}_{D_2/\mathbb{Q}_p}(1)^{-1}$$

in

$$\mathcal{M}^\times(\mathcal{O}_{C_{\mathbb{Q}_p}}(E), \Sigma(D_1) - \Sigma(D_2)) = \left(\text{Norm}_{D_1/\mathbb{Q}_p} \mathcal{O}_{C_{\mathbb{Q}_p}}(E) \otimes \text{Norm}_{D_2/\mathbb{Q}_p} \mathcal{O}_{C_{\mathbb{Q}_p}}(-E) \right)^\times$$

is sent to

$$\psi(\lambda \cdot \text{Norm}_{D_1/\mathbb{Q}_p}(1) \otimes \text{Norm}_{D_2/\mathbb{Q}_p}(1)^{-1}) := h_p(D_1 - D_2, E) + \log \lambda.$$

That this depends only on the linear equivalence classes of $D_1 - D_2$ and E follows from (6.4.4), plus (see [9, Proposition 5.2]) the fact that h_p is biadditive, symmetric and, for any non-zero rational function f on $C_{\mathbb{Q}_p}$ and any D in $\text{Div}^0(C_{\mathbb{Q}_p})$ with support disjoint from that of $\text{div}(f)$, we have $h_p(D, \text{div}(f)) = \log(f(D))$. Moreover, expressing h_p in terms of a Green function G as in [7, Theorem 7.3], we deduce that, in each residue disk of $\mathcal{M}^\times(\mathbb{Z}_p)$, ψ is given by a power series. Let $\omega_1, \dots, \omega_g$ be a basis of $\Omega_{C_{\mathbb{Q}_p}/\mathbb{Q}_p}^1(C_{\mathbb{Q}_p})$. This basis gives a unique morphism of groups $\log_J: J(\mathbb{Q}_p) \rightarrow \mathbb{Q}_p^g$ that extends the logarithm of Lemma 5.1.1. We define

$$\Psi := (\log_J \circ \text{opr}_{J,1}, \log_J \circ \text{opr}_{J,2}, \psi): \mathcal{M}^\times(\mathbb{Q}_p) \longrightarrow \mathbb{Q}_p^g \times \mathbb{Q}_p^g \times \mathbb{Q}_p.$$

By the biadditivity of h_p , Ψ is a morphism of biextensions, with the trivial biextension structure on $\mathbb{Q}_p^g \times \mathbb{Q}_p^g \times \mathbb{Q}_p$ as in (9.1.1). As $p > 2$, Ψ induces, from each residue polydisk to its image, a homeomorphism given by power series. Pulling back the coordinate functions on \mathbb{Q}_p^{2g+1} gives, for every $x \in \mathcal{M}^\times(\mathbb{F}_p)$, coordinates on $\mathcal{M}^\times(\mathbb{Z}_p)_x$.

We describe \tilde{j}_b and κ in these coordinates. It suffices to describe, for each i in $\{1, \dots, \rho-1\}$, $\tilde{j}_{b,i}: C \rightarrow T_i$, and from now on we omit the dependence on i . For each $c \in C(\mathbb{F}_p)$, on $T(\mathbb{Z}_p)_{\tilde{j}_b(x)}$ we use the coordinates $x_1 := f^*t_1, \dots, x_g := f^*t_g, z := f^*t_{2g+1}$ where f is the map $T \rightarrow \mathcal{M}^\times$ and t_1, \dots, t_{2g+1} are the coordinates on $\mathcal{M}^\times(\mathbb{Z}_p)_{\tilde{j}_b(c)}$ we just defined. Since the map Ψ is a morphism of biextensions, for j in $\{1, \dots, g\}$, $x_j \circ \kappa$ is a polynomial of degree at most 1, and $z \circ \kappa$ is a polynomial of degree at most 2. As explained in Section 7, over \mathbb{Z}_p , \tilde{j}_b is given by a line bundle \mathcal{L} over $(C \times C)_{\mathbb{Z}_p}$ rigidified along $(C \times \{b\})_{\mathbb{Z}_p}$ and along the diagonal with two sections l_b and l . Choosing a section that trivializes \mathcal{L} on an open subset of $(C \times C)_{\mathbb{Z}_p}$ containing (b, b) , (c, b) , and (c, c) in $(C \times C)(\mathbb{F}_p)$ we get a divisor D on $(C \times C)_{\mathbb{Z}_p}$ whose support is disjoint from (c, b) and (c, c) , and an isomorphism between \mathcal{L} and $\mathcal{O}(D)$ on $(C \times C)_{\mathbb{Z}_p}$. After modifying D with a principal horizontal divisor and a principal vertical divisor $D|_{C \times \{b\}}$ and $\text{diag}^* D$ are both equal to the zero divisor on $C_{\mathbb{Z}_p}$, hence l_b and l are the extensions of elements of \mathbb{Q}_p , interpreted as rational sections of $\mathcal{O}(D)$ on $(C \times C)_{\mathbb{Z}_p}$. By (7.5) and (7.8), there exists a unique $\lambda \in \mathbb{Q}_p^\times$ such that, for each $d \in C(\mathbb{Z}_p)_c$,

$$\tilde{j}_b(d) = \lambda \cdot \text{Norm}_{d/\mathbb{Z}_p}(1) \otimes \text{Norm}_{b/\mathbb{Z}_p}(1)^{-1} \in \mathcal{M}^\times(j_b(d), D|_{\{d\} \times C}).$$

Since x_j is the j -th coordinate of \log_J and since z is the pullback of ψ , we deduce that

$$x_1(\tilde{j}_b(d)) = \int_b^d \omega_1, \dots, x_g(\tilde{j}_b(d)) = \int_b^d \omega_g, \quad z(\tilde{j}_b(d)) = h_p(d-b, D|_{\{d\} \times C}) + \log \lambda,$$

and, by [3, Proof of Theorem 1.2] and [5, Lemma 5.5], the function $d \mapsto h_p(d-b, D|_{\{d\} \times C})$ is a sum of double Coleman integrals.

It should now be easy to exactly interpret geometrically the cohomological approach, showing that in the coordinates used here, the equations for $C(\mathbb{Q}_p)_2$ are precisely equations for the intersection of $C(\mathbb{Q}_p)$ and the p -adic closure of $T(\mathbb{Z})$. For doing computations, one can do them in the geometric context of this article, or, as in [5], in terms of the étale fundamental group of C . The connection between these is then given by p -adic local systems on T .

9.4 Work in progress, and future plans

We mention some work in progress and some research plans based on our geometric approach.

9.4.1 Generalisation to arbitrary number fields

At the Arizona Winter School, a group of 4 participants (Pavel Coupek, David Liliendfeldt, Luciana Xiao and Zijian Yao) started working on generalising the geometric approach from \mathbb{Q} to arbitrary number fields. In 4 days, they figured out how to generalise Theorem 4.12 and reported on it, see [10]. For generalisation of the cohomological approach, see [2] (quadratic Chabauty) and [11] (non-abelian Chabauty).

9.4.2 Modular curves

With Nicolas Mascot, Pietro Mercuri, and Pim Spelier, we are implementing our approach for finding all rational points on more modular curves such as $X(p)_{\text{ns}}^+$ and $X_0(p)^+$ with p prime

and not too large. We extend Khuri-Makdisi's methods in [18] for computing in jacobians (as used by Mascot in [20] and by Spelier in [25]) to Poincaré torsors, and work with these modular curves without using equations, but using modular forms in stead. We expect that this approach will outperform the cohomological approach. See for example the complexity estimates for the linear case in [25], Section 6.5. Pierre Parent is completing the determination of regular models of the $X(p)_{\text{ns}}^+$ over \mathbb{Z}_p .

9.4.3 Arakelov theory

The geometric approach is very close to Arakelov theory. Theorem 5.4 of [21] shows how P is naturally equipped with metrics on its fibres over $J(\mathbb{C}) \times J^\vee(\mathbb{C})$, and that for (x, y) in $(J \times J^{\vee,0})(\mathbb{Z})$, the Arakelov degree of $(x, y)^*P$ is the Néron-Tate height of (x, y) . Now T is the product of $\rho-1$ \mathbb{G}_m -torsors T_i on J . Their associated line bundles \mathcal{L}_i are pullbacks of P and so have natural metrics over $J(\mathbb{C})$. The pullbacks of the \mathcal{L}_i to U are trivial (uniquely up to signs). The squares of the norms of the trivialising sections are functions $g_i: U(\mathbb{C}) \rightarrow \mathbb{R}_{>0}$ with $(2\pi i)^{-1} \partial \bar{\partial} \log g_i$ equal to the pullback by $(\text{id}, m \cdot \circ \text{tr}_{c_i} \circ f_i)$ of the curvature form of the metric of P on $J(\mathbb{C}) \times J^\vee(\mathbb{C})$. The knowledge of $(2\pi i)^{-1} \partial \bar{\partial} \log g_i$ together with the value of g_i at b determine the g_i (see (7.8) and (7.2)). For any u in $U(\mathbb{Z})$ the Néron-Tate height of $(j_b(u), m(c_i + f_i(j_b(u))))$ is equal to the Arakelov height of $j_b(u)$ with respect to \mathcal{L}_i with its metric, and to the Arakelov degree of $(j_b(u), m(c_i + f_i(j_b(u))))^*P$, and the trivialising section makes this Arakelov degree equal to $(-1/2) \log g_i(u)$. This gives a bound on the absolute value of the Néron-Tate height of $(j_b(u), m(c_i + f_i(j_b(u))))$. This might help in determining the x in $J(\mathbb{Q})$ that are in $U(\mathbb{Z})$, but note that the functions $J(\mathbb{Q}) \rightarrow \mathbb{R}$ that send x to the Néron-Tate height of $(x, m(c_i + f_i(x)))$ are not definite. We intend to look at this in more detail. Especially the case where $r = \rho = g$ seems interesting to us.

9.4.4 Deeper quotients of the fundamental group

We have seen in Section 9.1 what quotient of $\pi_1(C(\mathbb{C}))$ is used in this article, it is a central extension of $\pi_1(C(\mathbb{C}))^{\text{ab}}$ by a quotient of $H_2(J(\mathbb{C}), \mathbb{Z})/H_2(C(\mathbb{C}), \mathbb{Z})$. We intend to see if there is a geometric interpretation for working with the extension by all of $H_2(J(\mathbb{C}), \mathbb{Z})/H_2(C(\mathbb{C}), \mathbb{Z})$.

Author contributions This project started with an idea of Edixhoven in December 2017. From then on Edixhoven and Lido worked together on the project. Section 8 is due entirely to Lido. Section 9 was written in July and August 2020.

Acknowledgements We thank Steffen Müller, Netan Dogra, Jennifer Balakrishnan, Kamal Khuri-Makdisi, Jan Vonk, Barry Mazur and Gerd Faltings for discussions and correspondence we had with them, Michael Stoll for suggesting the title of this article, Pim Spelier for his MSc thesis [25], and Sachi Hashimoto for the cartoon guide [15].

References

- [1] M.F. Atiyah, I.G. Macdonald, *Introduction to commutative algebra*. Addison–Wesley Publishing Co., Reading, Mass.–London–Don Mills, Ont. 1969
- [2] J. Balakrishnan, A. Besser, F. Bianchi, J. Steffen Müller, *Explicit quadratic Chabauty over number fields*. <https://arxiv.org/abs/1910.04653>
- [3] J. Balakrishnan, N. Dogra, *Quadratic Chabauty and rational points, I: p -adic heights*. With an appendix by J. Steffen Müller. *Duke Math. J.* 167 (2018), no. 11, 1981–2038.
- [4] J. Balakrishnan, N. Dogra, *An effective Chabauty–Kim theorem*. *Compos. Math.* 155 (2019), no. 6, 1057–1075.
- [5] J. Balakrishnan, N. Dogra, J.S. Müller, J. Tuitman, J. Vonk, *Explicit Chabauty–Kim for the split Cartan modular curve of level 13*. *Ann. of Math. (2)* 189 (2019), no. 3, 885–944.
- [6] D. Bertrand, B. Edixhoven, *Pink’s conjecture on unlikely intersections and families of semi-abelian varieties*. <https://arxiv.org/abs/1904.01788>
- [7] A. Besser, *p -adic Arakelov theory*. *J. Number Theory*, 111 (2005), no. 2, 318–371.
- [8] S. Bosch, W. Lütkebohmert, M. Raynaud, *Néron models*. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)* 21. Springer-Verlag, Berlin, 1990.
- [9] R. Coleman, G. Gross, *p -adic heights on curves*. *Algebraic number theory*, 7381 *Adv. Stud. Pure Math.*, 17 (1989).
- [10] P. Coupek, D. Lilienfeldt, L. Xiao, Z. Yao, *Geometric quadratic Chabauty over number fields*. <http://www.math.mcgill.ca/lilien/Chabauty-Part1.pdf>
- [11] N. Dogra, *Unlikely intersections and the Chabauty–Kim method over number fields*. <https://arxiv.org/abs/1903.05032>
- [12] B. Edixhoven, *Geometric quadratic Chabauty*. *Lectures at the Arizona Winter School 2020*. <http://swc.math.arizona.edu/index.html>
- [13] G. Faltings. *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*. *Invent. Math.* 73 (1983), no. 3, 349–366.
- [14] Y. Hasegawa, *Table of quotient curves of modular curves $X_0(N)$ with genus 2*. *Proc. Japan Acad. Ser. A Math. Sci.* 71 (1995), no. 10, 235–239 (1996).
- [15] S. Hashimoto, *Cartoon guide to finding \mathbb{Q} -points with geometric quadratic Chabauty*. <https://github.com/sachihashimoto/cartoon-guide-gqc>
- [16] T. Honda, *On the theory of commutative formal groups*. *Journal of the Mathematical Society of Japan* Vol. 22 No. 2 (1970). <https://projecteuclid.org/euclid.jmsj/1259942752>

- [17] N. Katz and B. Mazur, *Arithmetic moduli of elliptic curves*. Annals of Mathematics Studies, 108. Princeton University Press, Princeton, NJ, 1985.
- [18] K. Khuri-Makdisi, *Asymptotically fast group operations on Jacobians of general curves*. Math. Comp. 76 (2007), no. 260, 2213–2239.
- [19] Q. Liu, *Algebraic geometry and arithmetic curves*. Oxford Graduate Texts in Mathematics, 6. Oxford Science Publications. Oxford University Press, Oxford, 2002.
- [20] N. Mascot, *Hensel-lifting torsion points on Jacobians and Galois representations*. Math. Comp. 89 (2020), no. 323, 1417–1455.
- [21] L. Moret-Bailly, *Métriques permises*. Seminar on arithmetic bundles: the Mordell conjecture (Paris, 1983/84). Astérisque No. 127 (1985), 29–87.
http://www.numdam.org/article/AST_1985__127__29_0.pdf
- [22] L. Moret-Bailly, *Pinceaux de variétés abéliennes*. Astérisque No. 129 (1985).
http://www.numdam.org/item/AST_1985__129__1_0/
- [23] M. Raynaud, *Spécialisation du foncteur de Picard*. Inst. Hautes Études Sci. Publ. Math. No. 38 1970, 27-76. http://www.numdam.org/article/PMIHES_1970__38__27_0.pdf
- [24] *Groupes de monodromie en géométrie algébrique. I*. Séminaire de Géométrie Algébrique du Bois-Marie 1967–1969 (SGA 7 I). Dirigé par A. Grothendieck. Avec la collaboration de M. Raynaud et D.S. Rim. Lecture Notes in Mathematics, Vol 288. Springer–Verlag, Berlin–New York, 1972.
- [25] P. Spelier, *A geometric approach to linear Chabauty*. MSc thesis, Universiteit Leiden, 2020.
<https://www.universiteitleiden.nl/en/science/mathematics/education/theses>